

# **Geometria Algébrica I**

**lecture 5: Noetherian rings and irreducible affine varieties**

Misha Verbitsky

**IMPA, sala 232**

**August 31, 2018**

## Smooth points

**DEFINITION:** Let  $A \subset \mathbb{C}^n$  is an algebraic subset. A point  $a \in A$  is called **smooth**, if there exists a neighbourhood  $U$  of  $a \in \mathbb{C}^n$  (in usual topology) such that  $A \cap U$  is a smooth  $2k$ -dimensional real submanifold. A point is called **singular** if such diffeomorphism does not exist. A variety is called **smooth** if it has no singularities, and **singular** otherwise.

**PROPOSITION:** For any algebraic variety  $A$  and any smooth point  $a \in A$ , a diffeomorphism between a neighbourhood of  $a$  and an open ball **can be chosen polynomial**.

**Proof. Step1:** If  $a \in A \subset \mathbb{C}^n$  is a smooth point of a  $k$ -dimensional embedded manifold, **there exists  $k$  complex linear functions on  $\mathbb{C}^n$  such that their differentials are independent in the tangent space  $T_a A$ .**

**Step 2:** By inverse function theorem, these functions **define a diffeomorphism from a neighbourhood of  $A$  to an open subset of  $\mathbb{C}^k$ .** ■

## Analytic functions (reminder)

**REMARK:** In the coordinates defined by linear functions **all regular (polynomial) functions on  $A$  are analytic** (equal to the sum of their Taylor series).

In other polynomial coordinate systems, the Taylor series may be no longer finite, but **the regular functions remain analytic**, because the inverse function theorem remains true in analytic category.

**REMARK:** In fact, **all complex differentiable functions are analytic** (Cauchy), and the regular functions are clearly complex differentiable.

## Irreducible varieties

**DEFINITION:** An affine variety  $A$  is called **reducible** if it can be expressed as a union  $A = A_1 \cup A_2$  of affine varieties, such that  $A_1 \not\subset A_2$  and  $A_2 \not\subset A_1$ . If such a decomposition is impossible,  $A$  is called **irreducible**.

**CLAIM:** An affine variety  $A$  is **irreducible** if and only if its ring of polynomial functions  $\mathcal{O}_A$  **has no zero divisors**.

**Proof:** If  $A = A_1 \cup A_2$  is a decomposition of  $A$  into a non-trivial union of subvarieties, choose a non-zero function  $f \in \mathcal{O}_A$  vanishing at  $A_1$  and  $g$  vanishing at  $A_2$ . The product of these non-zero functions vanishes in  $A = A_1 \cup A_2$ , **hence**  $fg = 0$  **in**  $\mathcal{O}_A$ . Conversely, **if**  $fg = 0$ , **we decompose**  $A = V_f \cup V_g$ . ■

## Irreducibility for smooth varieties

**CLAIM:** Let  $M$  be an algebraic variety which is smooth and connected. **Then it is irreducible.**

**Proof:** (“Analytic continuation principle”). **Step 1**

Let  $f, g$  be non-zero polynomial functions,  $fg = 0$ . Decomposing  $f, g$  onto Taylor series around  $m \in M$ , **we obtain that the Taylor series for  $f$  or for  $g$  vanish.** Suppose it is  $f$  which has vanishing Taylor series, and let  $U \subset M$  be the set where all derivatives of  $f$  vanish.

**Step 2:** Since  $U$  is an intersection of closed sets  $\{x \in M \mid f^{(i)}(x) = 0\}$ , it is closed. However, an analytic function which has vanishing Taylor series in  $x$  has to vanish in a neighbourhood of  $x$ , hence  $U$  is also open. **An open and closed subset of  $M$  is  $M$  or  $\emptyset$ , because  $M$  is connected. ■**

## Irreducibility for smooth varieties (2)

**COROLLARY:** Let  $A$  be an affine variety such that its set  $A_0$  of smooth points is dense in  $A$  and connected. **Then  $A$  is irreducible.**

**Proof:** If  $f$  and  $g$  are non-zero functions such that  $fg = 0$ , the ring of polynomial functions on  $A_0$  contains zero divisors. However, **on a smooth, connected complex manifold the ring of polynomial functions has no zero divisors by analytic continuation principle.** ■

**REMARK:** Converse is also true: **an algebraic variety over  $\mathbb{C}$  is irreducible if and only if the set of its smooth points is connected.** This is a complicated result.

**EXERCISE:** Let  $X \rightarrow Y$  be a morphism of affine manifolds, where  $X$  is irreducible, and its image in  $Y$  is dense. **Prove that  $Y$  is also irreducible.**

## Noetherian rings and irreducible components

**DEFINITION:** A ring is called **Noetherian** if any increasing chain of ideals stabilizes: for any chain  $I_1 \subset I_2 \subset I_3 \subset \dots$  one has  $I_n = I_{n+1} = I_{n+2} = \dots$

**THEOREM: (Hilbert basis theorem)**

**Any finitely generated ring is Noetherian.**

**Proof:** *Later today.*

**DEFINITION: An irreducible component** of an algebraic set  $A$  is an irreducible algebraic subset  $A' \subset A$  such that  $A = A' \cup A''$ , and  $A' \not\subset A''$ .

**Remark 1:** Let  $A_1 \supset A_2 \supset \dots \supset A_n \supset \dots$  be a decreasing chain of algebraic subsets in an algebraic variety. **Then the corresponding ideals form an increasing chain of ideals:**  $\text{Ann}(A_1) \subset \text{Ann}(A_2) \subset \text{Ann}(A_3) \subset \dots$

**THEOREM:** Let  $A$  be an affine variety, and  $\mathcal{O}_A$  its ring of polynomial functions. Assume that  $\mathcal{O}_A$  is Noetherian. Then  **$A$  is a union of its irreducible components, which are finitely many.**

**Proof:** See the next slide ■

**Remark 2:** From the noetherianity and Remark 1 it follows that  **$A$  cannot contain a strictly decreasing infinite chain of algebraic subvarieties.**

## Noetherian rings and irreducible components (2)

**THEOREM:** Let  $A$  be an affine variety, and  $\mathcal{O}_A$  its ring of polynomial functions. Assume that  $\mathcal{O}_A$  is Noetherian. Then  **$A$  is a union of its irreducible components, which are finitely many.**

**Proof. Step 1:** Each point  $a \in A$  belongs to a certain irreducible component. Indeed, suppose that such a component does not exist. Then for each decomposition  $A = A_1 \cup A_2$  of  $A$  onto algebraic sets, the set  $A_i$  containing  $a$  can be split non-trivially onto a union of algebraic sets, the component containing  $a$  can also be split, and so on, *ad infinitum*. This gives a strictly decreasing infinite sequence, a contradiction (Remark 2).

**Step 2:** We proved existence of an irreducible decomposition, and **it remains only to show that number of irreducible components of  $A$  is finite.** Let  $A = \bigcup A_i$  be an irreducible decomposition.

**Step 3:** Let **algebraic closure** of a set  $X \subset \mathbb{C}^n$  be the intersection of all algebraic subsets of  $\mathbb{C}^n$  containing  $X$ . An intersection of algebraic sets is algebraic, because it is defined by the ideal generated by the union of their ideals. Since  $A = A_i \cup \bigcup_{j \neq i} A_j$ , the algebraic closure  $B_i$  of  $A \setminus A_i$  does not contain  $A_i$ . and **the sequence  $B_1 \supset B_1 \cap B_2 \supset B_1 \cap B_2 \cap B_3 \subset \dots$  decreases strictly, unless there are only finitely many irreducible components.** Applying Remark 2 again, we obtain that the number of  $B_i$  is finite. ■



## Noetherian rings

**DEFINITION:** A finitely generated ring is a quotient of a polynomial ring.

**THEOREM: (Hilbert's Basis Theorem)**

Any finitely generated ring over a field is Noetherian.

**Proof:** Later in this lecture.

**COROLLARY:** For any affine manifold, its ring of functions is Noetherian, hence the the irreducible decomposition exists and is finite.

**REMARK:** It suffices to prove Hilbert's Basis Theorem for the ring of polynomials. Indeed, any finitely generated ring is a quotient of the polynomial ring, but the set of ideals of the quotient ring  $A/I$  is injectively mapped to the set of ideals of  $R$ .

**REMARK:** Therefore, Hilbert's Basis Theorem would follow if we prove that  $R[t]$  is Noetherian for any Noetherian ring  $R$ .

**EXERCISE:** Find an example of a ring which is not Noetherian.

## Finitely generated ideals

**DEFINITION: Finitely generated ideal** in a ring is an ideal  $\langle a_1, \dots, a_n \rangle$  of sums  $\sum b_i a_i$ , where  $\{a_i\}$  is a fixed finite set of elements of  $R$ , called **generators** of  $R$ .

**LEMMA:** Let  $I \subset R$  be a finitely generated ideal, and  $I_0 \subset I_1 \subset I_2 \subset \dots$  an increasing chain of ideals, such that  $\bigcup_n I_n = I$ . **Then this chain stabilises.**

**Proof:** Let  $I = \langle a_1, \dots, a_n \rangle$ , and  $I_N$  be an ideal in the chain  $I_0 \subset I_1 \subset I_2 \subset \dots$  which contains all  $a_i$ . Then  $I_N = I$ . ■

**CLAIM: A ring  $R$  is Noetherian if and only if all its ideals are finitely generated.**

**Proof:** For any chain of ideals  $I_0 \subset I_1 \subset I_2 \subset \dots$ , **finite generatedness of  $I = \bigcup I_i$  guarantees stabilization of this chain**, as follows from Lemma above.

Conversely, if  $R$  is Noetherian, and  $I$  any ideal, take  $I_0 = 0$  and let  $I_k \subset I$  be obtained by adding to  $I_{k-1}$  an element of  $I$  not containing in  $I_{k-1}$ . **Since the chain  $\{I_k\}$  stabilizes,  $I$  is finitely generated.** ■

## Noetherian modules

**DEFINITION:** A module over a ring  $R$  is a vector space  $M$  equipped with an algebra homomorphism  $R \rightarrow \text{End}(M)$ .

**EXAMPLE:** A subspace  $I \subset R$  in a ring is an ideal if and only if  $I$  is an  $R$ -submodule of  $R$ , considered as an  $R$ -module.

**DEFINITION:** A module  $M$  over  $R$  is called **Noetherian** if any increasing chain of submodules of  $M$  stabilizes.

**REMARK:** Any submodules and quotient modules of a Noetherian  $R$ -module are again Noetherian.

## Finitely generated $R$ -modules

**DEFINITION:** An  $R$ -module is called **finitely generated** if it is a quotient of a **free module**  $R^n$  by its submodule.

**EXERCISE:** Show that **a module  $M$  is Noetherian iff any  $M' \subset M$  is finitely generated.** Use this to prove that **direct sums of Noetherian modules are Noetherian.**

**LEMMA: A ring  $R$  is Noetherian if and only if it is Noetherian as an  $R$ -module.**

**Proof:** Ideals in  $R$  is the same as  $R$ -submodules of  $R$ , stabilization of a chain of  $R$ -submodules in  $R$  is literally the same as stabilization of a chain of ideals in  $R$ . ■

**REMARK:** Let  $M$  be a module over  $R[t]$  which is Noetherian as an  $R$ -module, **Then it is Noetherian as  $R[t]$ -module.** ■

**COROLLARY:** If  $R$  is Noetherian, then  $R[t]/(t^N) = R^N$  is a Noetherian  $R$ -module. Therefore, **the ring  $R[t]/(t^N)$  is Noetherian.** ■

## Proof of Hilbert's basis theorem

**PROPOSITION:** Let  $R$  be a Noetherian ring. **Then the polynomial ring  $R[t]$  is also Noetherian.**

**Proof. Step 1:** Let  $I \subset R[t]$  be an ideal. We need to show that it is finitely generated. Consider the ideal  $I_0 \subset R$  generated by all leading coefficients of all  $P(t) \in I$ . Since  $R$  is Noetherian,  $I_0$  is finitely generated:  $I_0 = \langle a_1, \dots, a_n \rangle$ , where all  $a_i$  are leading coefficients of  $P_i(t) \in I$ .

**Step 2:** Let  $N$  be the maximum of all degrees of  $P_i$ . For each  $Q(t) \in I$  with the leading coefficient  $\sum a_i b_i$  **there exists a polynomial  $P_Q(t)$  of degree no bigger than  $N$  with the same leading coefficient:**  $P_Q(t) = \sum_i P_i(t) b_i t^{N - \deg P_i}$ .

**Step 3:** Let  $\tilde{Q}(t)$  be the remainder of the long division of  $Q(t) \in I$  by  $P_Q(t)$ . Then  $\tilde{Q}(t) = Q(t) \bmod \langle P_1(t), \dots, P_n(t) \rangle$ , and  $\deg \tilde{Q}(t) < N$ .

**Step 4:** We have constructed an  $R$ -module embedding

$$M := I / \langle P_1(t), \dots, P_n(t) \rangle \longrightarrow R[t] / (t^N).$$

Since  $M$  is a submodule of  $R[t] / (t^N)$ , it is a Noetherian module, as shown above, hence finitely generated. Pick a set of polynomials  $Q_1(t), \dots, Q_m(t) \in I$ , generating  $M$ . **Then  $\{Q_i(t), P_i(t)\}$  generate  $I$ . ■**