

Geometria Algébrica I

lecture 8: Tensor product of rings and fibered product

Misha Verbitsky

IMPA, sala 232

September 14, 2018

Tensor product (reminder)

DEFINITION: Let R be a ring, and M, M' modules over R . We denote by $M \otimes_R M'$ an R -module generated by symbols $m \otimes m'$, $m \in M, m' \in M'$, modulo relations

$$r(m \otimes m') = (rm) \otimes m' = m \otimes (rm'),$$

$$(m + m_1) \otimes m' = m \otimes m' + m_1 \otimes m',$$

$m \otimes (m' + m'_1) = m \otimes m' + m \otimes m'_1$ for all $r \in R, m, m_1 \in M, m', m'_1 \in M'$. Such an R -module is called **the tensor product of M and M' over R** .

REMARK: Suppose that M is generated over R by a set $\{m_i \in M\}$, and M' generated by $\{m'_j \in M'\}$. **Then $M \otimes_R M'$ is generated by $\{m_i \otimes m'_j\}$.**

THEOREM: (Universal property of the tensor product)

For any bilinear map $B : M_1 \times M_2 \rightarrow M$ **there exists a unique homomorphism $b : M_1 \otimes M_2 \rightarrow M$, making the following diagram commutative:**

$$\begin{array}{ccc} M_1 \times M_2 & \xrightarrow{B} & M_1 \otimes M_2 \\ & \searrow \gamma & \downarrow b \\ & & M \end{array}$$

■

Exactness of the tensor product (reminder)

THEOREM: Let $M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$ be an exact sequence of R -modules.
Then the sequence

$$M_1 \otimes_R M \longrightarrow M_2 \otimes_R M \longrightarrow M_3 \otimes_R M \longrightarrow 0 \quad (*)$$

is exact.

COROLLARY: Let $I \subset R$ be an ideal in a ring. **Then** $M \otimes_R (R/I) = M/IM$.

Proof: Apply the functor $\otimes_R M$ to the exact sequence $0 \longrightarrow I \longrightarrow R \longrightarrow R/I \longrightarrow 0$.
We obtain $IM \longrightarrow M \longrightarrow (R/I) \otimes_R M \longrightarrow 0$. ■

Tensor product of rings

DEFINITION: Let A, B be rings, $C \rightarrow A$, $C \rightarrow B$ homomorphisms. Consider A and B as C -modules, and let $A \otimes_C B$ be their tensor product. Define the ring multiplication on $A \otimes_C B$ as $a \otimes b \cdot a' \otimes b' = aa' \otimes bb'$. This defines **tensor product of rings**.

EXAMPLE: $\mathbb{C}[t_1, \dots, t_k] \otimes_{\mathbb{C}} \mathbb{C}[z_1, \dots, z_n] = \mathbb{C}[t_1, \dots, t_k, z_1, \dots, z_n]$. Indeed, if we denote by $\mathbb{C}_d[t_1, \dots, t_k]$ the space of polynomials of degree d , then $\mathbb{C}_d[t_1, \dots, t_k] \otimes_{\mathbb{C}} \mathbb{C}_{d'}[z_1, \dots, z_n]$ is polynomials of degree d in $\{t_i\}$ and d' in $\{z_i\}$.

EXAMPLE: For any homomorphism $\varphi : \mathbb{C} \rightarrow A$, **the ring $A \otimes_{\mathbb{C}} (C/I)$ is a quotient of A by the ideal $A \cdot \varphi(I)$** . This follows from $M \otimes_R (R/I) = M/IM$.

PROPOSITION: (associativity of \otimes)

Let $C \rightarrow A, C \rightarrow B, C' \rightarrow B, C' \rightarrow D$ be ring homomorphisms. Then $(A \otimes_C B) \otimes_{C'} D = A \otimes_C (B \otimes_{C'} D)$.

Proof: Universal property of \otimes implies that $\text{Hom}((A \otimes_C B) \otimes_{C'} D, M) = \text{Hom}(A \otimes_C (B \otimes_{C'} D), M)$ is the space of polylinear maps $A \otimes B \otimes D \rightarrow M$ satisfying $\varphi(ca, b, d) = \varphi(a, cb, d)$ and $\varphi(a, c'b, d) = \varphi(a, b, c'd)$. However, an object X of category is defined by the functor $\text{Hom}(X, \cdot)$ uniquely **(prove it)**.

■

Tensor product of rings and preimage of a point

DEFINITION: Recall that **the spectrum** of a finitely generated ring R is the corresponding algebraic variety, denoted by $\text{Spec}(R)$

PROPOSITION: Let $f : X \rightarrow Y$ be a morphism of affine varieties, $f^* : \mathcal{O}_Y \rightarrow \mathcal{O}_X$ the corresponding ring homomorphism, $y \in Y$ a point, and \mathfrak{m}_y its maximal ideal. **Denote by R_1 the quotient of $R := \mathcal{O}_X \otimes_{\mathcal{O}_Y} (\mathcal{O}_Y/\mathfrak{m}_y)$ by its nilradical. Then $\text{Spec}(R_1) = f^{-1}(y)$.**

Proof. Step 1: If $\alpha \in \mathcal{O}_Y$ vanishes in y , $f^*(\alpha)$ vanishes in all points of $f^{-1}(y)$. This implies that **the set V_I of common zeros of the ideal $I := \mathcal{O}_X \cdot f^*\mathfrak{m}_y$ contains $f^{-1}(y)$.**

Step 2: If $f(x) \neq y$, take a function $\beta \in \mathcal{O}_Y$ vanishing in y and non-zero in $f(x)$. Since $\varphi^*(\beta)(x) \neq 0$ and $\beta(y) = 0$, this gives $x \notin V_I$. **We proved that the set of common zeros of the ideal $I = \mathcal{O}_X \cdot f^*\mathfrak{m}_y$ is equal to $f^{-1}(y)$.**

Step 3: Now, strong Nullstellensatz implies that $\mathcal{O}_{f^{-1}(y)}$ is a quotient of $R = \mathcal{O}_X/I$ by nilradical. ■

Tensor product of rings and product of varieties

LEMMA: $A \otimes_C B \otimes_B B' = A \otimes_C B'$.

Proof: Follows from associativity of tensor product and $B \otimes_B B' = B'$. ■

LEMMA: $A \otimes_C (B/I) = A \otimes_C B / (1 \otimes I)$, where $1 \otimes I$ denotes the ideal $A \otimes_C I$.

Proof: Using $M \otimes_R (R/I) = M/IM$, we obtain

$$A \otimes_C (B/I) = (A \otimes_C B) \otimes_B (B/I) = (A \otimes_C B) / (1 \otimes I) \quad \blacksquare$$

Lemma 1: Let A, B be finitely generated rings without nilpotents, $R := A \otimes_C B$, and $N \subset R$ nilradical. **Then $\text{Spec}(R/N) = \text{Spec}(A) \times \text{Spec}(B)$.**

Proof. Step 1: Let $A = \mathbb{C}[t_1, \dots, t_n]/I$, $B = \mathbb{C}[z_1, \dots, z_k]/J$. Then $\mathbb{C}[t_1, \dots, t_n] \otimes_C \mathbb{C}[z_1, \dots, z_k] = \mathbb{C}[t_1, \dots, t_n, z_1, \dots, z_k]$. Applying the previous lemma twice, **we obtain $A \otimes_C B = \mathbb{C}[t_1, \dots, t_n, z_1, \dots, z_k]/(I + J)$** . Here $I + J$ means $I \otimes 1 \oplus 1 \otimes J$.

Step 2: The set V_{I+J} of common zeros of $I + J$ is $\text{Spec}(A) \times \text{Spec}(B) \subset \mathbb{C}^n \times \mathbb{C}^k$.

Step 3: Hilbert Nullstellensatz implies $\text{Spec}(R/N) = V_{I+J} = \text{Spec}(A) \times \text{Spec}(B)$. ■

REMARK: We shall see that **a tensor product $R := A \otimes_C B$ of reduced rings is reduced.**

Tensor product of rings and product of varieties (2)

LEMMA: For any finitely-generated ring A over \mathbb{C} , **intersection P of all its maximal ideals is its nilradical.**

REMARK: Let A, B be finite generated rings over \mathbb{C} , $B \rightarrow A$ a homomorphism, and $\mathfrak{m} \subset B$ a maximal ideal. Then the ring $A \otimes_B (B/\mathfrak{m})$ **can contain nilpotents**, even if A and B have no zero divisors.

EXERCISE: Give an example of such rings A, B .

THEOREM: Let A, B be finitely-generated, reduced rings over \mathbb{C} , and $R := A \otimes_{\mathbb{C}} B$ their product. **Then R is reduced** (that is, has no nilpotents).

Proof: see the next slide.

COROLLARY: $\text{Spec}(A) \times \text{Spec}(B) = \text{Spec}(A \otimes_{\mathbb{C}} B)$.

Tensor product of rings and product of varieties (2)

THEOREM: Let A, B be finitely-generated, reduced rings over \mathbb{C} , and $R := A \otimes_{\mathbb{C}} B$ their product. **Then R is reduced.**

Proof. Step1: By the previous lemma, it suffices to show that the **intersection P of maximal ideals of R is 0.**

Step 2: Let X, Y denote the varieties $\text{Spec}(A), \text{Spec}(B)$. Lemma 1 implies that **maximal ideals of R are points of $X \times Y$.**

Step 3: Every such ideal is given as $\mathfrak{m}_x \otimes \mathcal{O}_Y + \mathcal{O}_X \otimes \mathfrak{m}_y$, where $x \in X, y \in Y$. Then

$$P = \bigcap_{X \times Y} (\mathfrak{m}_x \otimes \mathcal{O}_Y + \mathcal{O}_X \otimes \mathfrak{m}_y) = \bigcap_Y \left(\left(\bigcap_X \mathfrak{m}_x \otimes \mathcal{O}_Y \right) + \mathcal{O}_X \otimes \mathfrak{m}_y \right) = \bigcap_Y \mathcal{O}_X \otimes \mathfrak{m}_y = 0.$$

This follows from $\bigcap_Y 1 \otimes \mathfrak{m}_y = \bigcap_X \mathfrak{m}_x \otimes 1 = 0$ since A and B are reduced. ■

Preimage and diagonal

Claim 2: Let $f : X \rightarrow Y$ be a morphism of algebraic varieties, $f^* : \mathcal{O}_Y \rightarrow \mathcal{O}_X$ the corresponding ring homomorphism, $Z \subset Y$ a subvariety, and I_Z its ideal. Denote by R_1 the quotient of a ring $R := \mathcal{O}_X \otimes_{\mathcal{O}_Y} (\mathcal{O}_Y/I_Z) = \mathcal{O}_X/f^*(I_Z)$ by its nilradical. **Then $\text{Spec}(R_1) = f^{-1}(Z)$.**

Proof: Clearly, the set of common zeros of the ideal $J := f^*(I_Z)$ contains $f^{-1}(Z)$. On the other hand, for any point $x \in X$ such that $f(x) \notin Z$ there exist a function $g \in J$ such that $g(x) \neq 0$. Therefore, $f^{-1}(Z) = V_J$, and strong Nullstellensatz implies that $\mathcal{O}_{f^{-1}(Z)} = R_1$. ■

Claim 3: Let M be an algebraic variety, and $\Delta \subset M \times M$ the diagonal, and $I \subset \mathcal{O}_M \otimes_{\mathbb{C}} \mathcal{O}_M$ the ideal generated by $r \otimes 1 - 1 \otimes r$ for all $r \in \mathcal{O}_M$. **Then \mathcal{O}_Δ is $\mathcal{O}_M \otimes_{\mathbb{C}} \mathcal{O}_M/I$.**

Proof. Step1: By definition of the tensor product, $\mathcal{O}_M \otimes_{\mathbb{C}} \mathcal{O}_M/I = \mathcal{O}_M \otimes_{\mathcal{O}_M} \mathcal{O}_M = \mathcal{O}_M$, hence it is reduced. If we prove that $\Delta = V_I$, the statement of the claim would follow from strong Nullstellensatz.

Step 2: Clearly, $\Delta \subset V_I$. To prove the converse, let $(m, m') \in M \times M$ be a point not on diagonal, and $f \in \mathcal{O}_M$ a function which satisfies $f(m) = 0, f(m') \neq 0$. Then $f \otimes 1 - 1 \otimes f$ is non-zero on (m, m') . ■

Fibered product

DEFINITION: Let $X \xrightarrow{\pi_X} M, Y \xrightarrow{\pi_Y} M$ be maps of sets. **Fibered product** $X \times_M Y$ is the set of all pairs $(x, y) \in X \times Y$ such that $\pi_X(x) = \pi_Y(y)$.

CLAIM: Let $X \xrightarrow{\pi_X} M, Y \xrightarrow{\pi_Y} M$ be morphism of algebraic varieties, $R := \mathcal{O}_X \otimes_{\mathcal{O}_M} \mathcal{O}_Y$, and R_1 the quotient of R by its nilradical. **Then $\text{Spec}(R_1) = X \times_M Y$.**

Proof: Let I be the ideal of diagonal in $\mathcal{O}_M \otimes_{\mathbb{C}} \mathcal{O}_M$. Since I is generated by $r \otimes 1 - 1 \otimes r$ (Claim 3), $R = \mathcal{O}_X \otimes_{\mathbb{C}} \mathcal{O}_Y / (\pi_X \times \pi_Y)^*(I)$. Applying Claim 2, we obtain that $\text{Spec}(R_1) = (\pi_X \times \pi_Y)^{-1}(\Delta)$. ■

Initial and terminal objects


DEFINITION: Commutative diagram in category \mathcal{C} is given by the following data. There is a directed graph (graph with arrows). For each vertex of this graph we have an object of category \mathcal{C} , and each arrow corresponds to a morphism of the associated objects. **These morphisms are compatible, in the following way.** Whenever there exist two ways of going from one vertex to another, the compositions of the corresponding arrows are equal.

DEFINITION: An initial object of a category is an object $I \in \mathcal{Ob}(\mathcal{C})$ such that $\text{Mor}(I, X)$ is always a set of one element. **A terminal object** is $T \in \mathcal{Ob}(\mathcal{C})$ such that $\text{Mor}(X, T)$ is always a set of one element.

EXERCISE: Prove that **the initial and the terminal object is unique.**

Limits and colimits of diagrams

DEFINITION: Let $S = \{X_i, \varphi_{ij}\}$ be a commutative diagram in \mathcal{C} , and $\vec{\mathcal{C}}_S$ be a category of pairs (object X in \mathcal{C} , morphisms $\psi_i : X \rightarrow X_i$, defined for all X_i) making the diagram formed by $(X, X_i, \psi_i, \varphi_{ij})$ commutative.



Morphisms $\text{Mor}(\{X, \psi_i\}, \{X', \psi'_i\})$, are morphisms $\Psi \in \text{Mor}(X, X')$, making the diagram formed by $(X, X', \psi_i, \psi'_i, \varphi_{ij})$ commutative. The terminal object in this category is called **limit**, or **inverse limit** of the diagram S .

DEFINITION: **Colimit**, or **direct limit** is obtained from the previous definition by inverting all arrows and replacing “terminal” by “initial”.

Products and coproducts

EXAMPLE: Let S be a diagram with two vertices X_1 and X_2 and no arrows. The inverse limit of S is called **product** of X_1 and X_2 , and inverse limit **the coproduct**.

EXAMPLE: Products in the category of sets, vector spaces and topological spaces are the usual products of sets, vector spaces and topological spaces (**check this**).

EXAMPLE: Coproduct in the category of groups is called **free product**, or **amalgamated product**. Coproduct of the group \mathbb{Z} with itself is called **free group**. Coproduct in the category of vector spaces is also the usual product of vector spaces.

Products and coproducts (2)

EXERCISE: Prove that **the product of algebraic varieties is their product in this category.**

EXERCISE: Prove that **coproduct of rings over \mathbb{C} in the category of rings is their tensor product.**


EXERCISE: Prove that **coproduct of reduced rings over \mathbb{C} in the category of reduced rings is the quotient of their tensor product over a nilradical.**

Since the category of algebraic varieties is equivalent to the category of finitely generated reduced rings, this gives another proof of the theorem.


THEOREM: Let A, B be finitely generated reduced rings over \mathbb{C} . **Then** $\text{Spec}(A \otimes_{\mathbb{C}} B/I) = \text{Spec}(A) \times \text{Spec}(B)$, where I is nilradical.

Fibered product

DEFINITION: Consider the following diagram:



Its limit is called **fibered product** of A and B over C . Colimit of the diagram



is called **coproduct** of A and B over C .

EXERCISE: Prove that the **fibered product of algebraic varieties is the same as their product in the category of algebraic varieties.**

EXERCISE: Prove that the **coproduct of rings A and B over C is $A \otimes_C B$.** Prove that the **coproduct of reduced rings A and B over C in the category of reduced rings is $A \otimes_C B/I$, where I is nilradical.**

Using strong Nullstellensatz again, we obtain

CLAIM: Let $X \xrightarrow{\pi_X} M, Y \xrightarrow{\pi_Y} M$ be morphisms of affine varieties, $R := \mathcal{O}_X \otimes_{\mathcal{O}_M} \mathcal{O}_Y$, and R_1 the quotient of R by its nilradical. **Then $\text{Spec}(R_1) = X \times_M Y$.**