

Geometria Algébrica I

lecture 14: Finite quotients and branched covers

Misha Verbitsky

IMPA, sala 232

October 5, 2018

Finite morphisms (reminder)

REMARK: Let M be a finitely generated R -module, and $R \rightarrow R'$ a ring homomorphism. **Then $M \otimes_R R'$ is a finitely generated R' -module.** Indeed, **if M is generated by x_1, \dots, x_n , then $M \otimes_R R'$ is generated by x_1, \dots, x_n .**

DEFINITION: A morphism $X \rightarrow Y$ of affine varieties is called **finite** if the ring \mathcal{O}_X is a finitely generated module over \mathcal{O}_Y . In this case, \mathcal{O}_X is called **an integral extension** of \mathcal{O}_Y .

THEOREM: Let $X \xrightarrow{f} Y$ be a finite morphism. Then for any point $y \in Y$, **the preimage $f^{-1}(y)$ is finite.**

Proof. Step 1: Since \mathcal{O}_X is finite generated as an \mathcal{O}_Y -module, the ring $R := \mathcal{O}_X \otimes_{\mathcal{O}_Y} (\mathcal{O}_Y/\mathfrak{m}_y)$ is finitely generated as an $\mathcal{O}_Y/\mathfrak{m}_y$ -module. Since $\mathcal{O}_Y/\mathfrak{m}_y = \mathbb{C}$, we obtain that R is an Artinian algebra over \mathbb{C} .

Step 2: Let $N \subset R$ be a nilradical. As shown in Lecture 9, **$\text{Spec}(R/N)$ is a finite set.**

Step 3: As shown in Lecture 8, **$\text{Spec}(R/N) = f^{-1}(y)$.** ■

Dominant, finite morphisms are surjective (reminder)

THEOREM: Let $f : X \rightarrow Y$ be a finite, dominant morphism of affine varieties. **Then f is surjective.**

Proof. Step 1: Restricting to irreducible components, we can always assume that Y , and hence X is irreducible. Let $A = \mathcal{O}_Y$, $B = \mathcal{O}_X$. We can consider A as a subring of B , which has no zero divisors, and assume that B is finitely generated as A -module.

Step 2: Let $\mathfrak{m}_y \subset A$ be a maximal ideal corresponding to $y \in Y$. **Nakayama's lemma implies that $\mathfrak{m}_y B \neq B$.**

Step 3: $f^{-1}(y) = \text{Spec}(B \otimes_A A/\mathfrak{m}_y) = \text{Spec}(B/\mathfrak{m}_y B)$. **Since this is non-zero ring, the set $f^{-1}(y)$ is non-empty. ■**

Finite quotients

CLAIM: Let R be a Noetherian ring without zero divisors, G a group acting by automorphisms on R , and R^G the ring of G -invariants. **Then $\varphi : \text{Spec } R \rightarrow \text{Spec } R^G$ is a finite, dominant morphism.**

Proof. Step 1: For any $g \in G$, consider the corresponding polynomial map $P_g : R \rightarrow R$, and let $r \in R$. The polynomial $P(t) := \prod_{g \in G} (t - g(r))$ has G -invariant coefficients for any $r \in R$, hence $P(t) \in R^G[t]$

Step 2: The morphism φ is finite because each $r \in R$ satisfies the equation $P(r) = 0$, where $P(t) = \prod_{g \in G} (t - g(r))$. It is dominant, because $R^G \subset R$. ■

DEFINITION: Let G be a finite group acting on an affine variety X by automorphisms. **The quotient space** X/G is $\text{Spec}(\mathcal{O}_X^G)$.

EXAMPLE: $\mathbb{C}^2/\{\pm 1\} = \mathbb{C}[x^2, y^2, xy] = \mathbb{C}[t_1, t_2, t_3]/(t_1 t_2 = t_3^2)$. Indeed, $\mathbb{C}^2/\{\pm 1\} = \text{Spec } A$, where $A = \mathbb{C}[x, y]^{\{\pm 1\}}$: **A is the ring of even polynomials.**

EXAMPLE: Let $G = \mathbb{Z}/n\mathbb{Z}$ act on \mathbb{C} by multiplication by a primitive root $\sqrt[n]{1}$. Then $\mathbb{C}/G = \text{Spec}(\mathbb{C}[t]^G) = \text{Spec}(\mathbb{C}[t^n])$, hence **the quotient space \mathbb{C}/G is isomorphic to \mathbb{C} .**

Finite quotients (2)

THEOREM: Consider the natural morphism $\text{Spec } R \xrightarrow{\varphi} \text{Spec } R^G$. Then $\varphi(x) = \varphi(y)$ if and only if $x \in G \cdot y$, that is, the **set of points in $\text{Spec } R^G$ is identified with the space of G -orbits.**

Proof. Step 1: If two maximal ideals of R are G -conjugated, their intersections of $R^G \subset R$ are equal. This gives $\varphi(gx) = \varphi(x)$: **each G -orbit is mapped to one point.** It remains to show that **the preimage of any point is exactly one G -orbit.**

Step 2: For any ideal $\mathfrak{m} \subset R^G$, one has $(\mathfrak{m}R)^G = \mathfrak{m}$ (lecture 6). Then $A^G = R^G/\mathfrak{m}$, where $A := R \otimes_{R^G} (R^G/\mathfrak{m}) = R/\mathfrak{m}R$.

Step 3: Let \mathfrak{m} be the maximal ideal of $y \in \text{Spec } R^G$, and N the nilradical of $A := R/\mathfrak{m}R$. Since $\varphi^{-1}(y) = \text{Spec}(A/N)$, **points of $\varphi^{-1}(y)$ are maximal ideals of the ring A/N .**

Step 4: A semisimple Artinian \mathbb{C} -algebra A/N is a direct sum of finite extensions of \mathbb{C} , which are all isomorphic to \mathbb{C} , giving $A/N = \bigoplus \mathbb{C}$. Since $A^G = \mathbb{C}$ (Step 2), **the group G acts on summands of $A/N = \bigoplus \mathbb{C}$ transitively.** Therefore, **all points of $\varphi^{-1}(y)$ belong to the same G -orbit. ■**

Transcendental extensions

CLAIM: Let $[k : \mathbb{C}]$ be an extension of \mathbb{C} , and $[K : k]$ an extension of k generated over k by $z \in K$. **Then either z is transcendental, and K is isomorphic to the field of rational functions $k(z)$, or z is algebraic, and $[K : k]$ is a finite extension.**

Proof: Indeed, either z is a root of polynomial, and then $[K : k]$ is finite, or K contains the polynomial ring $k[z]$, and then K contains $k(z)$. ■

LEMMA: Let $[K : k]$ be a finite extension, and $[K(t) : k(t)]$ the corresponding extension of rational functions. **Then $[K(t) : k(t)]$ is also finite.**

Proof: Primitive element theorem gives $K = k[\alpha]$, where α is an algebraic element, which is a root of a polynomial $P(z)$. Using the isomorphism $A/J \otimes_A B = B/JB$ (Lecture 7), we obtain

$$K(t) = K \otimes_k k(t) = \frac{k[z]}{(P(z))} \otimes_k k(t) = k(t)[z]/(P(z)).$$

■

EXERCISE: Find a proof which does not use existence of a primitive element.

Transcendence basis

DEFINITION: Let $k(t_1, \dots, t_n)$ be the field of rational functions of several variables, that is, the fraction field for the polynomial ring $k[t_1, \dots, t_n]$. Then the extension $[k(t_1, \dots, t_n) : k]$ is called **a purely transcendental extension of k** , and t_1, \dots, t_n are called **algebraically independent**.

REMARK: Clearly, t_1, \dots, t_n are algebraically independent if and only if there are no algebraic relations of form $P(t_1, \dots, t_n) = 0$, where P is a polynomial of n variables.

DEFINITION: **Transcendence basis** of an extension $[K : k]$ is a collection $z_1, \dots, z_n \in K$ generating a purely transcendental extension $K' := k(z_1, \dots, z_n)$ such that $[K : K']$ is an algebraic extension.

Transcendence basis in regular functions

Theorem 1: Let $k(X)$ be the field of rational functions on an irreducible affine variety X , with \mathcal{O}_X generated by t_1, \dots, t_n . Let $S \subset \{t_1, \dots, t_n\}$ be a maximal algebraically independent subset. **Then the extension $[k(X) : k(t_1, \dots, t_k)]$ is finite.**

Proof: Since \mathcal{O}_X is finitely generated, we can use induction by the number of generators t_1, \dots, t_n of \mathcal{O}_X . Let $A \subset \mathcal{O}_X$ be a subring generated by t_1, \dots, t_{n-1} , and t_1, \dots, t_k a transcendence basis on $k(A)$.

If t_n is algebraic over $k(A)$, then $[k(X) : k(A)]$ is finite; since $[k(A) : k(t_1, \dots, t_k)]$ is finite, this implies that $[k(X) : k(t_1, \dots, t_k)]$ is finite.

If t_n is transcendental over $k(A)$, we obtain $k(X) = k(A)(t_n)$, and $[k(X) : k(t_1, \dots, t_k)] = [k(A)(t_n) : k(t_1, \dots, t_k, t_n)]$ is finite over $k(t_1, \dots, t_k, t_n)$ by the lemma above. ■

Transcendence basis and dominant morphisms

PROPOSITION: Let $X \subset \mathbb{C}^n$ be an irreducible affine manifold, t_1, \dots, t_n coordinates on \mathbb{C}^n , and $\Pi_k : X \rightarrow \mathbb{C}^k$ the projection to the first k coordinates.

Then the following are equivalent.

(i) Π_k is dominant and the extension $[k(X) : k(t_1, \dots, t_k)]$ is finite.

(ii) t_1, \dots, t_k is transcendence basis in $k(X)$.

Proof: Theorem 1 implies that $[k(X) : k(t_1, \dots, t_k)]$ is finite whenever t_1, \dots, t_k is the transcendence basis. Therefore, (ii) \Rightarrow (i). Converse is clear, because $k(X)$ is algebraic over $k(t_1, \dots, t_k) \subset k(X)$, hence t_1, \dots, t_k is a maximal algebraically independent subset. ■

REMARK: We want to find a projection $\Pi_k : X \rightarrow \mathbb{C}^k$ which is finite, for any given irreducible affine variety X .

When the coordinate projection is finite

REMARK: Let $X \subset \mathbb{C}^n$ be an irreducible affine subvariety, z_i coordinates on \mathbb{C}^n , and z_1, \dots, z_k transcendence basis on $k(X)$. **The projection map Π_{n-1} is finite if and only if $P(z_n) = 0$ in \mathcal{O}_X , for some monic polynomial $P(t) \in \mathcal{O}_X[t]$ with coefficients which are polynomial in z_1, \dots, z_{n-1} .** Indeed, this is precisely what is needed for \mathcal{O}_X to be a finitely generated module over its subalgebra $A = \mathcal{O}_{\Pi_{n-1}(X)}$ generated by z_1, \dots, z_{n-1} considered as regular functions on X . Notice that **a non-zero polynomial with $P(t) \in A[t]$ with $P(z_n) = 0$ always exists, unless $n = k$ and $X = \mathbb{C}^n$,** but it is not necessarily monic.

CLAIM: In these assumptions, **there exists a linear coordinate change $z'_i := z_i + \lambda_i z_n$, such that z_n is finite over z'_1, \dots, z'_k .**

Proof. Step 1: Let $P(z_1, \dots, z_k, t)$ be a non-zero polynomial such that $P(z_1, \dots, z_k, z_n) = 0$ in \mathcal{O}_X . Such a polynomial exists because z_1, \dots, z_k is a transcendence basis in $k(X)$, and z_n is algebraic over $z_1, \dots, z_k \in \mathcal{O}_X$. Let $F(z_1, \dots, z_k, z_n)$ be a homogeneous component of maximal degree in $P(z_1, \dots, z_k, z_n)$. We choose P to be of minimal possible degree in z_1, \dots, z_k, z_n .

When the coordinate projection is finite (2)

CLAIM: In these assumptions, **there exists a linear coordinate change** $z'_i := z_i + \lambda_i z_n$, **such that z_n is finite over z'_1, \dots, z'_k .**

Proof. Step 1: Let $P(z_1, \dots, z_k, t)$ be a non-zero polynomial such that $P(z_1, \dots, z_k, z_n) = 0$ in \mathcal{O}_X . Such a polynomial exists because z_1, \dots, z_k is a transcendence basis in $k(X)$, and z_n is algebraic over $z_1, \dots, z_k \in \mathcal{O}_X$. Let $F(z_1, \dots, z_k, z_n)$ be a homogeneous component of maximal degree in $P(z_1, \dots, z_k, z_n)$. We choose P to be of minimal possible degree in z_1, \dots, z_k, z_n .

Step 2: Consider a polynomial

$$Q(z'_1, \dots, z'_k, z_n) := F(z_1 + \lambda_1 z_n, \dots, z_k + \lambda_k z_n, z_n).$$

Then $Q(0, 0, \dots, 0, 1) = F(\lambda_1, \dots, \lambda_k, 1)$ is non-zero on X for general λ_i . Indeed, if $F(\lambda_1, \dots, \lambda_k, 1)$ is identically 0 for all λ_i , the top degree homogeneous term in $P(z_1, \dots, z_k, z_n)$ vanishes on X , and we can replace $P(z_1, \dots, z_k, z_n)$ by a smaller degree polynomial.

Step 3: The degree d polynomial $Q(z'_1, \dots, z'_k, t)$ is monic in t , because its leading term t^d has non-zero coefficient by Step 2. ■

Noether's normalization lemma, first version

REMARK: Let $A \subset B \subset C$ be ring without zero divisors, C is finitely generated as B -module, and B as an A -module. Then **C is finitely generated as an A -module.**

REMARK: We have proven that the projection $\Pi_{n-1} : X \rightarrow \mathbb{C}^{n-1}$ to coordinates $z'_1, \dots, z'_k, z_{k+1}, \dots, z_{n-1}$ is finite. Using this remark and induction by n , we obtain

COROLLARY: (Noether's normalization lemma, first version)

Let $X \subset \mathbb{C}^n$ be an irreducible affine subvariety. Then there exists a linear coordinate change such that **the projection of X to the first k coordinates gives a finite map $X \rightarrow \mathbb{C}^k$.** ■