

Teoria Ergódica Diferenciável

lecture 8: Riemannian geometry of space forms

Instituto Nacional de Matemática Pura e Aplicada

Misha Verbitsky, October 6, 2017

Riemannian manifolds

DEFINITION: Let $h \in \text{Sym}^2 T^*M$ be a symmetric 2-form on a manifold which satisfies $h(x, x) > 0$ for any non-zero tangent vector x . Then h is called **Riemannian metric**, of **Riemannian structure**, and (M, h) **Riemannian manifold**.

DEFINITION: For any $x, y \in M$, and any piecewise smooth path $\gamma : [a, b] \rightarrow M$ connecting x and y , consider **the length** of γ defined as $L(\gamma) = \int_{\gamma} \left| \frac{d\gamma}{dt} \right| dt$, where $\left| \frac{d\gamma}{dt} \right| = h\left(\frac{d\gamma}{dt}, \frac{d\gamma}{dt}\right)^{1/2}$. Define **the geodesic distance** as $d(x, y) = \inf_{\gamma} L(\gamma)$, where infimum is taken for all paths connecting x and y .

EXERCISE: Prove that the **geodesic distance satisfies triangle inequality and defines a metric on M** .

EXERCISE: Prove that **this metric induces the standard topology on M** .

EXAMPLE: Let $M = \mathbb{R}^n$, $h = \sum_i dx_i^2$. **Prove that the geodesic distance coincides with $d(x, y) = |x - y|$** .

EXERCISE: Using partition of unity, **prove that any manifold admits a Riemannian structure**.

Conformal structures

DEFINITION: Let h, h' be Riemannian structures on M . These Riemannian structures are called **conformally equivalent** if $h' = fh$, where f is a positive smooth function.

DEFINITION: **Conformal structure** on M is a class of conformal equivalence of Riemannian metrics.

DEFINITION: **A Riemann surface** is a 2-dimensional oriented manifold equipped with a conformal structure.

Almost complex structures

DEFINITION: Let $I : TM \rightarrow TM$ be an endomorphism of a tangent bundle satisfying $I^2 = -\text{Id}$. Then I is called **almost complex structure operator**, and the pair (M, I) **an almost complex manifold**.

CLAIM: Let M be a 2-dimensional oriented conformal manifold. **Then M admits a unique orthogonal almost complex structure** in such a way that the pair $x, I(x)$ is positively oriented. Conversely, **an almost complex structure uniquely determines the conformal structure and orientation**.

Proof: The almost complex structure is $\frac{\pi}{2}$ degrees counterclockwise rotation; it is clearly determined by the conformal structure and orientation. To prove that the conformal structure is recovered from the almost complex structure, define the action of $U(1)$ on TM as follows: $\rho(t) = e^{tI}$. Any I -invariant metric is also ρ -invariant, hence constant on circles which are its orbits. Therefore all such metrics are proportional. ■

Homogeneous spaces

DEFINITION: A Lie group is a smooth manifold equipped with a group structure such that the group operations are smooth. Lie group G **acts on a manifold** M if the group action is given by the smooth map $G \times M \rightarrow M$.

DEFINITION: Let G be a Lie group acting on a manifold M transitively. Then M is called **a homogeneous space**. For any $x \in M$ the subgroup $\text{St}_x(G) = \{g \in G \mid g(x) = x\}$ is called **stabilizer of a point** x , or **isotropy subgroup**.

CLAIM: For any homogeneous manifold M with transitive action of G , **one has** $M = G/H$, where $H = \text{St}_x(G)$ is an isotropy subgroup.

Proof: The natural surjective map $G \rightarrow M$ putting g to $g(x)$ identifies M with the space of conjugacy classes G/H . ■

REMARK: Let $g(x) = y$. Then $\text{St}_x(G)^g = \text{St}_y(G)$: **all the isotropy groups are conjugate**.

Isotropy representation

DEFINITION: Let $M = G/H$ be a homogeneous space, $x \in M$ and $\text{St}_x(G)$ the corresponding stabilizer group. The **isotropy representation** is the natural action of $\text{St}_x(G)$ on T_xM .

DEFINITION: A Riemannian form Φ on a homogeneous manifold $M = G/H$ is called **invariant** if it is mapped to itself by all diffeomorphisms which come from $g \in G$.

REMARK: Let Φ_x be an isotropy invariant scalar product on T_xM . For any $y \in M$ obtained as $y = g(x)$, consider the form Φ_y on T_yM obtained as $\Phi_y := g(\Phi)$. The choice of g is not unique, however, for another $g' \in G$ which satisfies $g'(x) = y$, we have $g = g'h$ where $h \in \text{St}_x(G)$. Since Φ_x is h -invariant, **the metric Φ_y is independent from the choice of g .**

We proved

THEOREM: Homogeneous Riemannian forms on $M = G/H$ are in bijective correspondence with isotropy invariant spalar products on T_xM , for any $x \in M$. ■

Space forms

DEFINITION: **Simply connected space form** is a homogeneous manifold of one of the following types:

positive curvature: S^n (an n -dimensional sphere), equipped with an action of the group $SO(n+1)$ of rotations

zero curvature: \mathbb{R}^n (an n -dimensional Euclidean space), equipped with an action of isometries

negative curvature: $SO(1, n)/SO(n)$, equipped with the natural $SO(1, n)$ -action. This space is also called **hyperbolic space**, and in dimension 2 **hyperbolic plane** or **Poincaré plane** or **Bolyai-Lobachevsky plane**

Riemannian metric on space forms

LEMMA: Let $G = SO(n)$ act on \mathbb{R}^n in a natural way. **Then there exists a unique G -invariant symmetric 2-form:** the standard Euclidean metric.

Proof: Let g, g' be two G -invariant symmetric 2-forms. Since S^{n-1} is an orbit of G , we have $g(x, x) = g(y, y)$ for any $x, y \in S^{n-1}$. Multiplying g' by a constant, we may assume that $g(x, x) = g'(x, x)$ for any $x \in S^{n-1}$. **Then $g(\lambda x, \lambda x) = g'(\lambda x, \lambda x)$ for any $x \in S^{n-1}, \lambda \in \mathbb{R}$;** however, all vectors can be written as λx . ■

COROLLARY: Let $M = G/H$ be a simply connected space form. **Then M admits a unique, up to a constant multiplier, G -invariant Riemannian form.**

Proof: The isotropy group is $SO(n-1)$ in all three cases, and the previous lemma can be applied. ■

REMARK: From now on, all space forms are assumed to be homogeneous Riemannian manifolds.