

## K3 surfaces, assignment 7: Moser isotopy lemma

**Exercise 7.1.** Let  $M$  be a compact manifold, and  $V_0, V_1$  two smooth volume forms which satisfy  $\int_M V_0 = \int_M V_1$ . Prove that there exists a diffeomorphism which satisfies  $\Phi^*V_0 = V_1$ .

**Exercise 7.2.** Let  $(M, I)$  be an almost complex manifold, and  $\omega_0, \omega_1$  cohomologous symplectic forms which satisfy  $\omega_i(x, Ix) > 0$  for any non-zero tangent vector  $x$  (such forms are called **taming**).

- Prove that there exists a diffeomorphism  $\Phi$  which satisfies  $\Phi^*\omega_0 = \omega_1$ .
- Prove that  $|x|_i^2 := \omega_i(x, Ix)$  is a Hermitian metric on  $M$ . Prove that a diffeomorphism that satisfies  $\Phi^*\omega_0 = \omega_1$  defines an isometry

$$(M, |x|_1^2) \longrightarrow (M, |x|_0^2)$$

if  $\Phi$  is compatible with  $I$ , that is, satisfies  $d\Phi(Ix) = I(d\Phi(x))$ .

- Find examples of  $\omega_0, \omega_1$  taming  $(M, I)$  such that a diffeomorphism compatible with  $I$  and satisfying  $\Phi^*\omega_0 = \omega_1$  does not exist.

**Exercise 7.3.** Let  $\omega_i \in \Lambda^2(M)$  be a sequence of symplectic forms on a compact manifold  $M$  converging to a symplectic form  $\omega$  in  $C^0$ -topology. Assume that all  $\omega_i$  are homologous. Prove that almost all  $(M, \omega_i)$  are symplectomorphic.

**Exercise 7.4.** Let  $(M, V)$  be a connected manifold equipped with a volume form. Prove that the group of volume-preserving diffeomorphisms acts on  $M$  transitively.

**Exercise 7.5.** Prove that the group of symplectomorphisms of  $(M, \omega)$  acts transitively on  $M$ , for any connected symplectic manifold  $(M, \omega)$ .

**Exercise 7.6 (\*).** Let  $\Psi$  be a diffeomorphism of a connected symplectic manifold  $(M, \omega)$  which satisfies  $\Psi^*\omega = 2\omega$ . Prove that  $\omega$  is exact, or find a counterexample.