

K3 surfaces

lecture 26: Reflections in cohomology and Kähler chambers

Misha Verbitsky

IMPA, sala 236

December 2, 2024, 17:00

Complex structures on a K3 surface (reminder)

THEOREM: Let M be a complex manifold diffeomorphic to a K3 surface. **Then M is K3**, that is, satisfies $K_M = \mathcal{O}_M$.

Proof. Step 1: Since $b_1(M)$ is even, M is Kähler by Buchdahl-Lamari. Also, $c_2(M) = 24$ because it is its topological Euler characteristic. Since the signature of M is $(3, 19)$, Hodge index theorem implies $\dim H^{2,0}(M) = 1$. Now, $b_1(M) = 0$ implies that $\chi(\mathcal{O}_M) = 2$ and $\text{rk } H^0(K_M) = 1$. Riemann-Roch formula for surfaces gives $\chi(\mathcal{O}_M) = \frac{c_1^2 + c_2}{12} = \frac{c_1^2}{2} + 2$, hence $c_1^2 = 0$. This implies that $\chi(K_M^{\otimes i}) = 2$ for all i . **If we prove that $\text{rk } H^0(K_M^{\otimes i}) > 0$ for all i , it would imply that K_M is trivial.**

Step 2: Assume, by contradiction, that K_M is non-trivial. Since K_M is effective, $\text{rk } H^0(K_M^{\otimes i}) = 0$ for all $i < 0$. Serre's duality gives $\text{rk } H^0(K_M^{\otimes i}) = H^2(K_M^{\otimes -i+1})$ and $\text{rk } H^1(K_M^{\otimes i}) = \text{rk } H^1(K_M^{\otimes -i+1})$. Then $\chi(K_M^{\otimes i}) = 2$ implies that $H^0(K_M^{\otimes i}) \geq 2$ for all $i > 1$. The corresponding sections of $K_M^{\otimes i}$ don't intersect, because $c_1^2 = 0$, hence the line system $K_M^{\otimes i}$ is globally generated and defines a holomorphic map π to $\mathbb{C}P^1$. The bundle $K_M^{\otimes i}$ restricted to any fiber of π has degree 0, hence the fibers of π are 1-dimensional. **This implies that M is a surface of Kodaira dimension 1.**

Step 3: From Kodaira-Enriques classification it follows that all surfaces of Kodaira dimension 1 are elliptic with base a curve S of genus ≥ 1 . Therefore, **$\pi_1(M)$ surjects to $\pi_1(S)$ and $H_1(M)$ is infinite.** This brings a contradiction.

■

Teichmüller space for complex structures (reminder)

DEFINITION: Let Teich be the Teichmüller space of complex structures of Kähler type on a K3 surface. The corresponding period space is denoted $\mathbb{P}\text{er} := \{v \in \mathbb{P}H^2(M, \mathbb{C}) \mid \int_M v \wedge v = 0, \int_M v \wedge \bar{v} > 0\}$.

PROPOSITION: (local Torelli theorem)

Let Teich be the space of complex structures on a K3 surface, and $\mathbb{P}\text{er} : \text{Teich} \rightarrow \mathbb{P}\text{er}$ the map taking (M, I) to the line $H^{2,0}(M) \subset H^2(M, \mathbb{C})$. **Then $\mathbb{P}\text{er}$ is a local diffeomorphism.**

Proof: Lecture 17. ■

Kähler chambers (reminder)

DEFINITION: Let $V \in \mathbb{P}er = \text{Gr}_{++}(H^2(M, \mathbb{R}))$, and $I \in \text{Teich}$ be a complex structure such that $\text{Per}(I) = V$. The set $\text{Kah}(I) \subset \text{Pos}(I)$ is called **a Kähler chamber** of V . **The set of Kähler chambers** for V is the set of all $\text{Kah}(I)$ for all $I \in \text{Per}^{-1}(V)$.

THEOREM: Let $V \in \mathbb{P}er = \text{Gr}_{++}(H^2(M, \mathbb{R}))$, and \mathfrak{X}_V be the set of all (-2) -classes orthogonal to V . Let $S_V := \bigcup_{\eta \in \mathfrak{X}_V}$. Then the Kähler chambers of V are connected components of $\text{Pos}(V) \setminus S_V$.

Proof. Step 1: Let $I \in \text{Per}^{-1}(V)$ be a complex structure. **Its Kähler cone is the set of all classes $\omega \in \text{Pos}(V)$ which are positive on effective (-2) -classes;** clearly, this set is one of the connected components of $\text{Pos}(V) \setminus S_V$.

Step 2: Consider the forgetful map $\text{Teich}_H \longrightarrow H^2(M, \mathbb{R}) \times \text{Teich}$ taking (I, J, K, g) to (I, ω_I) . Since $\text{Teich}_H = \mathbb{P}er_H$, this map is surjective on the set of all $v \in \text{Pos}(V) \setminus S_V$. Therefore, **any of the connected components of $\text{Pos}(V) \setminus S_V$ is realized as a Kähler cone for some $I \in \text{Per}^{-1}(V)$.** ■

Global Torelli theorem for Teichmüller spaces (reminder)

THEOREM: Let Teich be a connected component of the Teichmüller space of all complex structures on a K3 surface, and $\text{Per} : \text{Teich} \rightarrow \mathbb{P}\text{er}$ the period map. **Then Per is surjective, and bijective for all $V \in \mathbb{P}\text{er} \subset \text{Gr}_{++}(H^2(M, \mathbb{R}))$ such that V^\perp does not contain (-2) -curves.**

Proof. Step 1: Consider the space $\mathbb{P}\text{er}_1$ of all pairs $\{(W \times V) \in \mathbb{P}\text{er}_h \times \mathbb{P}\text{er} \mid V \subset W\}$. This space is S^2 -fibered over $\mathbb{P}\text{er}_h$, with the fiber being the set of all oriented 2-planes in $W \in \mathbb{P}\text{er}_h$. Similarly, let Teich_1 be the set of all pairs $(\mathcal{H}, L) \in \text{Teich}_h \times \text{Teich}$, consisting of all hyperkähler structures $\langle I, J, K \rangle$ inducing L ; this space is also S^2 -fibered over Teich . This gives a commutative diagram:

$$\begin{array}{ccc} \text{Teich}_1 & \xrightarrow{P} & \text{Teich} \\ \downarrow \text{Per}_1 & & \downarrow \text{Per} \\ \mathbb{P}\text{er}_1 & \xrightarrow{\Psi} & \mathbb{P}\text{er} \end{array}$$

Step 2: Torelli theorem for $\mathbb{P}\text{er}_h$ implies that Per_1 is a diffeomorphism. Calabi-Yau theorem implies that the fiber $P^{-1}(I)$ of $\text{Teich}_1 \xrightarrow{P} \text{Teich}$ is projectivization of the Kähler cone $\text{Kah}(I)$ of (M, I) ; the fiber of $\mathbb{P}\text{er}_1 \xrightarrow{\Psi} \mathbb{P}\text{er}$ is the positive cone $\text{Pos}(I)$. Since Per_1 is a diffeomorphism, Ψ is one to one on points for which $\text{Kah}(I) = \text{Pos}(I)$; for other points, $\Psi^{-1}(V) = \mathfrak{K}_V$, where \mathfrak{K}_V is the set of Kähler chambers. ■

Automorphisms of K3 surfaces acting trivially on $H^2(M)$

Proposition 1: Let ν be an automorphism of a K3 surface (M, I) acting trivially on $H^2(M, \mathbb{Z})$. **Then $\nu = \text{Id}$.**

Proof. Step 1: By Calabi-Yau theorem, ν acts by hyperkähler isometries for any hyperkähler metric on (M, I) . Using the same argument as in the proof of Torelli theorem for hyperkähler structures, **we can extend ν to a hyperkähler isometry of any other hyperkähler structure in Teich_h .**

Step 2: Then ν acts by an automorphism of a Kummer surface $M = \widetilde{X}/\pm 1$, where X is any complex 2-torus. When X is very non-algebraic (e. g. contains no complex curves), it is easy to see that **$\text{Aut } M$ is identified with the group of $\{\pm 1\}$ -invariant automorphisms of X .** Since ν acts trivially on $H^2(X)$, it is induced by a translation of the torus. Finally, translations which are $\{\pm 1\}$ -invariant are translations by 2-torsion. The group G of such translations is clearly isomorphic to the group $(\mathbb{Z}/2)^4$. This group acts freely and transitively on 16 singular points of $X/\pm 1$. Therefore, all non-trivial elements of G act on $H^2(M)$ non-trivially. ■

Marked K3 surfaces

DEFINITION: A **marking** on a K3 surface M is an isomorphism $H^2(M, \mathbb{Z}) \cong \Lambda$, where $\Lambda = (-E_8)^2 \oplus (U_2)^3$ is the lattice of the second cohomology of M .

REMARK: Fix an isomorphism $H^2(M, \mathbb{Z}) \cong \Lambda$ on a smooth K3 surface, and let Teich be the corresponding Teichmüller space. Then each $I \in \text{Teich}$ corresponds to a complex structure on a marked K3 surface. Conversely, global Torelli theorem implies that **each marked K3 surface (M, I) is obtained from a point in a Teichmüller space.**

DEFINITION: Let Γ be the mapping class group of a K3 surface M . **The Torelli group** is the kernel of the natural map $\Gamma \rightarrow O(H^2(M, \mathbb{Z}))$.

PROPOSITION: **The space of marked K3 surfaces is naturally identified with the Teichmüller space of K3 surfaces.**

Proof: Since the quartics are dense in any connected component of Teichmüller space of K3 surfaces, the Torelli group acts transitively on the connected components of Teich . This action is also free, because any automorphism of a K3 acting trivially on $H^2(M)$ is trivial (Proposition 1). Therefore, a K3 surface with a marking uniquely determines a point in Teich . ■

Reflections in the mapping class group

Proposition 2: Let M be a K3 surface, and $v \in H^2(M, \mathbb{Z})$ a (-2) -class. Consider the corresponding reflection in cohomology, $r_v(x) := x + (r, x)r$, where $(r, x) = \int_M r \wedge x$ is the intersection form. Then there exists a family of K3 surfaces $\mathcal{M} \xrightarrow{\pi} X$ **such that its monodromy acts as r_v on $H^2(M)$.**

Proof. Step 1: Let $V \in \text{Gr}_{++}(H^2(M, \mathbb{R}))$ be a point in the period space such that $V^\perp \cap H^2(M, \mathbb{Z}) = \langle v \rangle$. A general positive plane in $\langle v \rangle^\perp$ has this property **(prove it)**. Let $W \subset H^2(M, \mathbb{R})$ be a 3-space $W = V + \langle v \rangle$. The set $\text{Gr}_{++}(W)$ is a complex analytic subset of $\text{Gr}_{++}(H^2(M, \mathbb{R}))$ biholomorphic to a disk Δ . **This defines a holomorphic family $V_t \in \text{Gr}_{++}(H^2(M, \mathbb{R}))$, $t \in \Delta$, which is invariant with respect to r_v .**

Step 2: Replacing Δ by a smaller disk, and using the local Torelli theorem, we can assume that there exists a family $\mathcal{M} \rightarrow \Delta$ of K3 surfaces over $\Delta \subset V_t$, with the fibers (M, I_t) over $V_t \in \Delta$. For all $t \in \Delta$ outside of a countable set Q , we have $V_t^\perp = 0$. By global Torelli theorem, this implies that **(M, I_t) is uniquely, up to an isomorphism, determined by V_t for all $t \in \Delta \setminus Q$.**

Step 3: Therefore, the action of r_v takes a complex structure I_t associated with $t \in \Delta \setminus Q$ to an isomorphic complex structure with different marking, defining a holomorphic family over the quotient space $\frac{\Delta \setminus Q}{\langle r_v \rangle}$. **This family has monodromy r_v .** ■

The mapping class group of a K3 surface

EXERCISE: Let M be a K3 surface, and $\mathcal{P} \subset H^2(M, \mathbb{R})$ the space of all positive real vectors. **Prove that \mathcal{P} is homotopy equivalent to S^2 .** Denote by $O^+(H^2(M, \mathbb{R}))$ the index 2 subgroup of $O(H^2(M, \mathbb{R}))$ of all isometries preserving the generator of $H^2(\mathcal{P}, \mathbb{Z})$. **Prove that $O^+(H^2(M, \mathbb{R}))$ is an index 2 subgroup of $O(H^2(M, \mathbb{R}))$.**

COROLLARY: (Borcea, Donaldson)

Let Mon be the image of the mapping class group of K3 in $O(H^2(M, \mathbb{Z}))$.

Then Mon is an index 2 subgroup obtained as the intersection $O(H^2(M, \mathbb{Z})) \cap O^+(H^2(M, \mathbb{R}))$.

Proof: As shown by C. T. C. Wall, $O^+(H^2(M, \mathbb{Z}))$ is an index 2 subgroup in $O(H^2(M, \mathbb{Z}))$ generated by reflections r_v for all (-2) -classes v (*C. T. C. Wall, On the orthogonal groups of unimodular quadratic forms II, J. Reine Angew. Math. 213. (1963/64), 122-136.*). Each r_v belongs to Mon as follows from Proposition 2 (this was the original contribution of Borcea). Finally, $\text{Mon} \subset O^+(H^2(M, \mathbb{Z}))$, as shown in *S. K. Donaldson, The orientation of Yang-Mills moduli spaces and 4-manifold topology, J. Diff. Geom. 26 (1987), 397-428.* ■

The Weyl group of a K3 surface

DEFINITION: Let (M, I) be a K3 surface, and $\mathfrak{R} \subset H^{1,1}(M, \mathbb{Z})$ the set of all (-2) -classes of type $(1,1)$. **The Weyl group** of a K3 is the group generated by the reflections r_v for all $v \in \mathfrak{R}$.

THEOREM: The Weyl group W of K3 surface **acts transitively on the set of Kähler chambers.**

Proof: Let Mon_I be the set of all $A \in O^+(H^2(M, \mathbb{Z}))$ preserving the Hodge structure on $H^2(M, \mathbb{R})$. From the explicit description of Kähler chambers it follows that Mon_I takes a Kähler chambers to a Kähler chambers. **Therefore, the Weyl group W takes a Kähler chamber to a Kähler chamber.**

Step 2: Let $K \subset \text{Pos}(M, I)$ be a Kähler chamber, and v a (-2) -vector such that v^\perp contains a face F of K . Then r_v takes K to a Kähler chamber adjacent to F . This way, one obtains every Kähler chamber adjacent to K from K . Iterating this construction, we obtain Kähler chambers adjacent to ones adjacent to K , and so on. **Therefore, $W \cdot K$ is the set of all Kähler chambers.** ■

COROLLARY: Let $I_1, I_2 \in \text{Teich}$ be two points with the same periods, $\text{Per}(I_1) = \text{Per}(I_2)$. **Then (M, I_1) and (M, I_2) is the same K3 with different markings.**

Proof: The action of each generator of the Weyl group changes the marking on the same K3 surface. ■

Hodge-theoretic Torelli theorem

DEFINITION: Let M be a K3 surface. **A K3-type Hodge structure** on $H^2(M)$ is a decomposition $H^2(M, \mathbb{C}) = H^{2,0}(M) \oplus H^{1,1}(M) \oplus H^{0,2}(M)$ such that $\overline{H^{p,q}(M)} = H^{q,p}(M)$, the intersection pairing between $H^{q,p}(M)$ and $H^{p_1,q_1}(M)$ vanishes unless $p = q_1, q = p_1$, and $\int_M \Omega \wedge \overline{\Omega} > 0$ for any non-zero $\Omega \in H^{2,0}(M)$.

THEOREM: Let M be a K3 surface, Γ its mapping class group, and \mathfrak{S} the set of all K3-type Hodge structures up to $O^+(H^2(M, \mathbb{Z}))$ -action. **Then the natural map from $\text{Teich} / \Gamma \rightarrow \mathfrak{S}$ is bijective.** In other words, **the K3-type Hodge structures are in bijective correspondence with classes of isomorphism of K3 surfaces.**

Proof: Clearly, $\mathfrak{S} = \frac{\mathbb{P}\text{er}}{O^+(H^2(M, \mathbb{Z}))}$, hence $\text{Per} : \text{Teich} \rightarrow \mathbb{P}\text{er}$ is reduced to $\text{Per} : \text{Teich} / \Gamma \rightarrow \frac{\mathbb{P}\text{er}}{O^+(H^2(M, \mathbb{Z}))}$. This map is surjective by global Torelli theorem. For any $V \in \mathbb{P}\text{er}$, the set $\text{Per}^{-1}(V)$ is identified with the set of Kähler chambers in $\text{Pos}(V^\perp)$. The Weyl group acts transitively on the set of Kähler chambers. Therefore, **the set $\text{Per}^{-1}(V)$ is mapped to the same point in Teich / Γ .**

Step 2: This implies that all $I \in \text{Teich}$ which are mapped to the same point in $\frac{\mathbb{P}\text{er}}{O^+(H^2(M, \mathbb{Z}))}$ are related by the Γ -action. ■

Automorphism group of a K3 surface

REMARK: Let (M, I) be a K3 surface. **Proposition 1 implies that the natural map $\text{Aut}(M, I) \longrightarrow O^+(H^2(M, \mathbb{Z}))$ is injective.** In the sequel, **we will identify $\text{Aut}(M, I)$ and its image in $O^+(H^2(M, \mathbb{Z}))$.**

THEOREM: Let (M, I) be a K3 surface. Then $\text{Aut}(M, I) \subset O^+(H^2(M, \mathbb{Z}))$ **is the group of all $\rho \in O^+(H^2(M, \mathbb{Z}))$ which preserve the Hodge decomposition and the Kähler cone of (M, I) .**

Proof. Step 1: Clearly, any $\rho \in \text{Aut}(M, I)$ preserves the Hodge decomposition and the Kähler cone. **It remains to prove the converse:** any $\rho \in O^+(H^2(M, \mathbb{Z}))$ preserving the the Hodge decomposition and the Kähler cone is induced by an automorphism of (M, I) .

Step 2: Consider the action of ρ on the space $\text{Teich}_H = \mathbb{P}er_H$ of hyperkähler structures (I, J, K, g) . This action is extended to the corresponding universal fibration, because points of Teich_H are hyperkähler structures on marked K3, and $O^+(H^2(M, \mathbb{Z}))$ just changes the marking.

Step 3: Let $\Psi : \text{Teich}_H \longrightarrow \text{Teich}$ take (I, J, K, g) to $I \in \text{Teich}$. Since ρ preserves the Kähler cone and the Hodge decomposition, it takes $\Psi^{-1}(I)$ to $\Psi^{-1}(I)$. Therefore, ρ takes (I, J, K, g) to (I, J', K', g') ; this map is by construction holomorphic. ■