

# **Complex manifolds of dimension 1**

**lecture 12: Tilings, hyperelliptic curves, Ananin theorem**

Misha Verbitsky

**IMPA, sala 232**


**February 10, 2020**

## Homotopy lifting principle (reminder)

**DEFINITION:** A topological space  $X$  is **locally path connected** if for each  $x \in X$  and each neighbourhood  $U \ni x$ , there exists a smaller neighbourhood  $W \ni x$  which is path connected.

## THEOREM: (homotopy lifting principle)

Let  $X$  be a simply connected, locally path connected topological space, and  $\tilde{M} \rightarrow M$  a covering map. Then for each continuous map  $X \rightarrow M$ , there exists a lifting  $X \rightarrow \tilde{M}$  making the following diagram commutative.



## Coverings and subgroups of $\pi_1(M)$

**THEOREM:** For each subgroup  $\Gamma \subset \pi_1(M)$  **there exists a unique, up to isomorphism, connected covering  $M_\Gamma \rightarrow M$  such that  $\pi_1(M_\Gamma) = \Gamma$ .**

**THEOREM:** If, in addition,  $\Gamma \subset \pi_1(M)$  is a normal subgroup, **the group  $G = \pi_1(M)/\Gamma$  acts on  $M_\Gamma$  by automorphisms** commuting with projection to  $M$  (“automorphisms of the covering”), freely and transitively on the fibers of the projection  $M_\Gamma \rightarrow M$ , and give  $M = M_\Gamma/G$ .

**COROLLARY:** The fundamental group  $\pi_1(M)$  **acts on the universal covering  $\tilde{M}$  by homeomorphisms which commute with the projection to  $M$  and give  $M = \tilde{M}/\pi_1(M)$ .**

**THEOREM:** Let  $M$  be connected, locally path connected, locally simply connected topological space. Fix  $p \in M$ . Then **the category of the coverings of  $M$  is naturally equivalent with the category of sets with the action of  $\pi_1(M)$** , and the equivalence takes a covering  $\tilde{M} \rightarrow M$  to the set  $\pi^{-1}(p)$ .

**COROLLARY:** Let  $M$  be a space with commutative  $\pi_1(M)$ , and  $\tilde{M}$  its universal cover. **Then for any connected covering  $M_1 \rightarrow M$ , the covering  $M_1$  is obtained as  $M_1 = \tilde{M}/\Gamma$ , where  $\Gamma \subset \pi_1(M)$  is a subgroup.**

## Deformation retracts

**DEFINITION: Retraction** of a topological space  $X$  to  $Y \subset X$  is a continuous map  $X \rightarrow Y$  which is identity on  $Y \subset X$ . **Deformation retraction** of a topological space  $X$  to  $Y \subset X$  is a continuous map  $\varphi_t : X \times [0, 1] \rightarrow X$  such that  $\varphi_1 = \text{Id}_X$  and  $\varphi_0$  is retraction of  $X$  to  $Y$ .

**EXERCISE: Prove that  $\pi_1(X) = \pi_1(Y)$  when  $Y$  is a deformation retract of  $X$ .**

**DEFINITION:** A topological space  $X$  is **contractible** if a point  $p \in X$  is its deformation retract.

**EXERCISE:** Let  $p \in X$  be a deformation retract of  $X$ , **prove that any other point  $q \in X$  is also a deformation retract.**

**EXERCISE:** Prove that **a contractible space  $X$  satisfies  $\pi_1(X) = 0$ .**

**EXERCISE:** Let  $Y \subset X$  be a deformation retract of  $X$ . **Prove that any map  $Z \rightarrow X$  is homotopy equivalent to  $Z \rightarrow Y \subset X$ .**

## Points of ramification

**DEFINITION:** Let  $\varphi : X \rightarrow Y$  be a holomorphic map of complex manifolds, not constant on each connected component of  $X$ . Any point  $x \in X$  where  $d\varphi = 0$  is called **a ramification point** of  $\varphi$ . **Ramification index** of the point  $x$  is the number of preimages of  $y' \in Y$ , for  $y'$  in a sufficiently small neighbourhood of  $y = \varphi(x)$ .

**THEOREM 1:** Let  $X, Y$  be compact Riemannian surfaces,  $\varphi : X \rightarrow Y$  a holomorphic map, and  $x \in X$  a ramification point. Then there is a neighbourhood of  $x \in X$  biholomorphic to a disk  $\Delta$ , **such that the map  $\varphi|_{\Delta}$  is equivalent to  $\varphi(x) = x^n$** , where  $n$  is the ramification index.

**Proof. Step 1:** Let  $W \subset Y$  be a sufficiently small simply connected neighbourhood of  $y \in Y$ , and  $U \ni x$  a connected component of its preimage in  $X$ . Choosing  $W$  sufficiently small, we may assume that  $U$  lies in a coordinate chart. The zeros of  $d\varphi$  are isolated. Shrinking  $W$  if necessary, we may assume that  $d\varphi$  is nowhere zero on  $U \setminus x$ , and  $U \setminus x \xrightarrow{\varphi} W \setminus y$  is a covering. We identify  $W$  with a disk  $\Delta$ . By homotopy lifting principle, the homothety map  $\lambda \rightarrow r\lambda$  of  $W$ ,  $r \in [0, 1]$  can be lifted to  $U$  uniquely. **This means that  $x$  is a homotopy retract of  $U$ , and  $\pi_1(U) = 0$ .** Riemann mapping theorem implies that  $U$  **is isomorphic to a disk.**

## Points of ramification (2)

**THEOREM 1:** Let  $X, Y$  be compact Riemannian surfaces,  $\varphi : X \rightarrow Y$  a holomorphic map, and  $x \in X$  a ramification point. Then there is a neighbourhood of  $x \in X$  biholomorphic to a disk  $\Delta$ , **such that the map  $\varphi|_{\Delta}$  is equivalent to  $\varphi(x) = x^n$** , where  $n$  is the ramification index.

**Proof. Step 1:** Let  $W \subset Y$  be a sufficiently small simply connected neighbourhood of  $y \in Y$ , and  $U \ni x$  a connected component of its preimage in  $X$ . [...] Choose  $U, W$  in such a way that that  $x$  is a homotopy retract of  $U$ , and  $\pi_1(U) = 0$ . **Riemann mapping theorem implies that  $U$  is isomorphic to a disk.**

**Step 2:** Passing to the universal covering  $\widetilde{U} \setminus x = \widetilde{W} \setminus y$ , we obtain an holomorphic action of  $\mathbb{Z} = \pi_1(\widetilde{U} \setminus x)$  on  $\widetilde{W} \setminus y$  such that  $W \setminus y = \widetilde{W} \setminus y / \mathbb{Z}$  and  $U \setminus y = \widetilde{W} \setminus y / n\mathbb{Z}$ . Therefore,  $\mathbb{Z}/n\mathbb{Z}$  acts on  $U \setminus x$ , freely and transitively on the fibers of the projection  $U \setminus x \xrightarrow{\varphi} W \setminus y$ . This action is extended to 0 by homotopy lifting principle. **Then  $W = U / (\mathbb{Z}/n)$ .** However, **any action of the cyclic group  $\mathbb{Z}/n$  on  $\Delta$  is conjugate to the rotations by  $\{\varepsilon_n^i\}$** , where  $\varepsilon_n$  is a primitive root of unity of degree  $n$ . The corresponding quotient map is equivalent to  $\varphi(x) = x^n$ . ■

## Hyperelliptic curves and hyperelliptic equations

**REMARK:** Let  $M \xrightarrow{\Psi} N$  be a  $C^\infty$ -map of compact smooth oriented manifolds. Recall that **degree** of  $\Psi$  is number of preimages of a regular value  $n$ , counted with orientation. Recall that **the number of preimages is independent from the choice of a regular value  $n \in N$** , and **the degree is a homotopy invariant**.

**DEFINITION: Hyperelliptic curve**  $S$  is a compact Riemann surface admitting a holomorphic map  $S \rightarrow \mathbb{C}P^1$  of degree 2 and with  $2n$  ramification points of degree 2.

**DEFINITION: Hyperelliptic equation** is an equation  $P(t, y) = y^2 + F(t) = 0$ , where  $F \in \mathbb{C}[t]$  is a polynomial with no multiple roots.

**REMARK:** Clearly, the natural projection  $(t, y) \rightarrow t$  maps the set  $S_0$  of solutions of  $P(t, y) = 0$  to  $\mathbb{C}$  with  $2n$  ramification points of degree 2. Also,  $S_0$  is smooth (**check this**). The complex manifold  $S_0$  **is equipped with an involution  $\tau(t, y) = (t, -y)$  exchanging the roots, and  $S_0/\tau = \mathbb{C}$** .

## Hyperelliptic curves and desingularization

**DEFINITION:** Let  $P(t, y) = y^2 + F(t) = 0$  be a hyperelliptic equation. **Homogeneous hyperelliptic equation** is  $P(x, y, z) = y^2 z^{n-2} + z^n F(x/z) = 0$ , where  $n = \deg F$ .

**REMARK:** The set of solutions of  $P(x, y, z) = 0$  is singular, but an algebraic variety of dimension 1 has a natural desingularization, called **normalization**. **The involution  $\tau$  is extended to the desingularization  $S$** , giving  $S/\tau = \mathbb{C}P^1$  because  $\mathbb{C}P^1$  is the only smooth holomorphic compactification of  $\mathbb{C}$  as we have seen already.



## Hyperbolic polyhedral manifolds

**DEFINITION:** A **polyhedral manifold** of dimension 2 is a piecewise smooth manifold obtained by gluing polygons along edges.

**DEFINITION:** Let  $\{P_i\}$  be a set of polygons on the same hyperbolic plane, and  $M$  be a polyhedral manifold obtained by gluing these polygons. Assume that all edges which are glued have the same length, and we glue the edges of the same length. Then  $M$  is called **a hyperbolic polyhedral manifold**. We consider  $M$  as a metric space, with the path metric induced from  $P_i$ .

**CLAIM:** Let  $M$  be a hyperbolic polyhedral manifold. Then for each point  $x \in M$  which is not a vertex,  **$x$  has a neighbourhood which is isometric to an open set of a hyperbolic plane.**

**Proof:** For interior points of  $M$  this is clear. When  $x$  belongs to an edge, it is obtained by gluing two polygons along isometric edges, hence the neighbourhood is locally isometric to the union of the same polygons in  $\mathbb{H}^2$  aligned along the edge. ■

## Hyperbolic polyhedral manifolds: interior angles of vertices

**DEFINITION:** Let  $v \in M$  be a vertex in a hyperbolic polyhedral manifold. **Interior angle** of  $v$  in  $M$  is sum of the adjacent angles of all polygons adjacent to  $v$ .

**EXAMPLE 1:** Let  $M \rightarrow \Delta$  be a ramified  $n$ -tuple cover of the Poincaré disk, given by solutions of  $y^n = x$ . We can lift split  $\Delta$  to polygons and lift the hyperbolic metric to  $M$ , obtaining  $M$  as a union of  $n$  times as many polygons glued along the same edges. **Then the interior angle of the ramification point is  $2\pi n$ .**

**EXAMPLE 2:** Let  $\Delta \rightarrow M$  be a ramified  $n$ -tuple cover, obtained as a quotient  $M = \Delta/G$ , where  $G = \mathbb{Z}/n\mathbb{Z}$ . Split  $\Delta$  onto fundamental domains of  $G$ , shaped like angles adjacent to 0. Then the quotient  $\Delta/G$  gives an angle with its opposite sides glued. **It is a hyperbolic polyhedral manifold with interior angle  $\frac{2\pi}{n}$  at its ramification point.**

**EXAMPLE 3:** Let  $D$  be a diameter bisecting a disk  $\Delta$ , and passing through the origin 0 and  $P \subset \Delta$  one of the halves. The (unique) edge of  $P$  is split onto two half-geodesics  $E_+$  and  $E_-$  by the origin. **Gluing  $E_+$  and  $E_-$ , we obtain a hyperbolic polyhedral manifold with a single vertex and the interior angle  $\pi$ .**

## Sphere with $n$ points of angle $\pi$

**EXAMPLE:** Let  $P$  be a bounded convex polygon in  $\mathbb{H}^2$ ,  $\alpha$  the sum of its angles, and  $a_i$ ,  $i = 1, \dots, m$  median points on its edges  $E_i$ . Each  $a_i$  splits  $E_i$  in two equal intervals. We glue them as in Example 3, and glue all vertices of  $P$  together. **This gives a sphere  $M$  with hyperbolic polyhedral metric**, one vertex  $\nu$  with angle  $\alpha$  (obtained by gluing all vertices of  $P$  together) and  $m$  vertices with angle  $\pi$  corresponding to  $a_i \in E_i$ .

**REMARK:** Assume that  $\alpha = 2\pi$ , that is,  $M$  is isometric to a hyperbolic sphere around  $\nu$ . We equip  $M$  with a complex structure compatible with the hyperbolic metric outside of its singularities. A neighbourhood of each singularity is isometrically identified with a neighbourhood of 0 in  $\Delta/G$ , where  $G = \mathbb{Z}/2\mathbb{Z}$ . We put a complex structure on  $\Delta/G$  as in Example 2. **This puts a structure of a complex manifold on  $M$ .**

### **THEOREM: (Alexandre Ananin)**

Let  $M$  be the hyperbolic polyhedral manifold obtained from the polyhedron  $P$  as above. Assume that  $m = n$  is even, and  $\alpha = 2\pi$ . **Then  $M$  admits a double cover  $M_1$ , ramified at all  $a_i$ , which is locally isometric to  $\mathbb{H}^2$ .**

*Proof in the next lecture.*

**REMARK:** Clearly,  $M_1$  is hyperelliptic.