

Complex surfaces

lecture 7 bis: Differential operators (supplementary lecture)

Misha Verbitsky

IMPA, sala 236

(distributed for your perusal)

Differential operators

Notation: Let M be a smooth manifold, TM its tangent bundle, $\Lambda^i M$ the bundle of differential i -forms, $C^\infty M$ the smooth functions. **The space of sections of a bundle B is denoted by B .**

DEFINITION: Let M be a manifold. The ring of **differential operators on the ring of functions on M** is a subalgebra of $\text{End}_{\mathbb{R}}(C^\infty M, C^\infty M)$ is defined as follows. **Operator of order 0** is a $C^\infty M$ -linear map, that is, a map $L_\alpha : f \mapsto \alpha f$, where $\alpha \in C^\infty M$ is a smooth function. **Operator of order 1** is a sum of a differentiation along a vector field and a $C^\infty M$ -linear map. **Differential operator of order k** is a linear combination of products of k first order differential operators.

REMARK: In coordinates x_1, \dots, x_n , differential operators can be expressed as sums of **differential monomials**:

$$D = f_0 + \sum_{i=1}^n f_i \frac{\partial}{\partial x_i} + \sum_{i,j=1}^n f_{ij} \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i,j,k=1}^n f_{ijk} \frac{\partial^3}{\partial x_i \partial x_j \partial x_k} + \dots$$

Differential operators with coefficients in a vector bundle

DEFINITION: Let E, F be trivial vector bundles on M , with basis e_1, \dots, e_n in E , f_1, \dots, f_m in F . **A differential operator from E to F** is a function mapping $\sum_{i=1}^n \alpha_i e_i$, where $\alpha_i \in C^\infty M$, to

$$D \left(\sum_{i=1}^n \alpha_i e_i \right) = \sum_{j=1}^m \sum_{i=1}^n D_{ij}(\alpha_i) f_j, \quad (*)$$

where D_{ij} are differential operators on $C^\infty M$. One can think of D as a **$n \times m$ -matrix with coefficients in differential operators on $C^\infty M$** .

DEFINITION: We say that a section b of a vector bundle B on M **has support in a set $K \subset M$** if b vanishes in an open set which contains $M \setminus K$. The smallest of all such K is called **support** of b .

DEFINITION: Let E, F be vector bundles on M . Let D be an operator mapping sections of E to sections of F . Suppose that for any open set $U \subset M$ such that E and F are trivial on U with bases $\{e_i\}, \{f_j\}$, and for any $e = \sum_{i=1}^n \alpha_i e_i$ with support in U , the section $D(e)$ is expressed as in (*):

$$D \left(\sum_{i=1}^n \alpha_i e_i \right) = \sum_{j=1}^m \sum_{i=1}^n D_{ij}(\alpha_i) f_j.$$

Then D is called **a differential operator from E to F** .

Associated graded rings

REMARK: Algebra is an associative ring over a field. Rings in this lecture are not necessarily commutative, but always associative.

DEFINITION: Let R be an associative ring. **Filtration** on R is a collection of subspaces $R_0 \subset R_1 \subset R_2 \subset \dots$ such that $R_i R_j \subset R_{i+j}$.

REMARK: Let $x \in R_k, y \in R_l$. Then the product xy modulo R_{k+l-1} depends only on the class of x modulo R_{k-1} . Indeed, $R_{k-1} R_l \subset R_{k+l-1}$. **This defines the product map** $(R_k/R_{k-1}) \otimes (R_l/R_{l-1}) \longrightarrow R_{k+l}/R_{k+l-1}$. **We obtained the associative product structure on the space** $\bigoplus_{i=0}^{\infty} R_i/R_{i-1}$.

DEFINITION: Let $R_0 \subset R_1 \subset R_2 \subset \dots$ be a filtered ring. The ring $\bigoplus_{i=0}^{\infty} R_i/R_{i-1}$ is called **the associated graded ring** of this filtration.

EXAMPLE: Consider the filtration $\text{Diff}^0(M) \subset \text{Diff}^1(M) \subset \text{Diff}^2(M) \subset \dots$ on the ring of differential operators. The associated graded ring is called **the ring of symbols of differential operators**.

Order of zeroes of a function

DEFINITION: Let $m \in M$, and x_1, \dots, x_n a coordinate system around m , with $x_1(m) = x_2(m) = \dots = x_n(m) = 0$. We say that **a function f has zero of order $\geq k$ at m** if $\frac{\partial^l f}{\partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_l}}(m) = 0$ for any $l < k$.

CLAIM: Let $\mathfrak{m} \subset C^\infty M$ be the ideal of all functions vanishing in $m \in M$. **Then f has zero of order $\geq k$ at m if and only if $f \in \mathfrak{m}^k$.**

Proof. Step 1: Let $f \in \mathfrak{m}^k$. Then $\frac{\partial f}{\partial x_i} \in \mathfrak{m}^{k-1}$ by Leibniz rule. **Hence f has zero of order $\geq k$ at m .**

Step 2: The function f has zero of order $\geq k$ at m if and only if $\frac{\partial f}{\partial x_i}$ has zero of order $\geq k - 1$ at m .

Step 3: If f has zero of order ≥ 1 at m , this means that $f \in \mathfrak{m}$ by definition. Using induction in k and step 2, we obtain that **f has zero of order $\geq k$ at $m \Leftrightarrow \frac{\partial f}{\partial x_i} \in \mathfrak{m}^{k-1} \Leftrightarrow f \in \mathfrak{m}^k$.** ■

COROLLARY: Consider a function f with the Taylor series decomposition $f = \sum P_i$ in m , where $P_i \in \mathbb{R}[x_1, \dots, x_n]$ a homogeneous polynomial of degree i . **Then f has zero of order $\geq k$ in m if and only if $P_0 = P_1 = \dots = P_{k-1} = 0$.**

■

Differential operators: algebraic definition

DEFINITION: (Grothendieck) Let R be a commutative ring over a field k . Given $a \in R$, consider the map $L_a : R \rightarrow R$ mapping x to ax . Define $\text{Diff}^k(R) \subset \text{Hom}_k(R, R)$ inductively as follows. The $\text{Diff}^0(R)$ is the space of all R -linear maps from R to R , that is, the space of all L_a , $a \in R$. The space $\text{Diff}^k(R)$, $k > 0$ is

$$\text{Diff}^k(R) := \{D \in \text{Hom}_k(R, R) \mid [L_a, D] \in \text{Diff}^{k-1}(R) \forall a \in R.\}$$

EXERCISE: Prove that $\text{Diff}^k(C^\infty M)$ in this sense is the same as the space of differential operators defined in the standard way.

CLAIM: For any $D \in \text{Diff}^{k-1}(C^\infty M)$, and any $a \in C^\infty M$, one has $[L_a, D] \in \text{Diff}^{k-1}(C^\infty M)$.

Proof: Indeed, for a vector field $X \in TM$, one has $[L_a, \text{Lie}_X] = L_{\text{Lie}_X(a)}$, which means that $[L_a, \text{Diff}^1(C^\infty M)] \subset \text{Diff}^0(C^\infty M)$. Now, if we take a commutator of L_a and a product of k elements from $\text{Diff}^1(C^\infty M)$, we obtain a linear combination of products of $k-1$ elements of $\text{Diff}^1(C^\infty M)$, by Leibniz formula:

$$[L_a, D_1 D_2 \dots D_n] = [[L_a, D_1] D_2 \dots D_n + D_1 [L_a, D_2] D_3 \dots D_n + \dots + D_1 D_2 \dots D_{n-1} [L_a, D_n]].$$

■

Differential operators and homogeneous polynomials

LEMMA: Let x_1, \dots, x_n be coordinates on \mathbb{R}^n , and $D \in \text{Diff}^k(\mathbb{R}^n)$ a differential operator vanishing on all homogeneous polynomials $P \in \mathbb{R}[x_1, \dots, x_n]$ of degree k . **Then $D = 0$.**

Proof. Step 1: We prove lemma by induction. For $k = 0$ it is clear. Let $L_{x_i} : C^\infty M \rightarrow C^\infty M$ be the multiplication by x_i . Then $[L_{x_i}, D]$ is a differential operator of order $k - 1$ vanishing on all homogeneous polynomials of degree $\leq k - 1$. Using induction on k , we obtain that $[L_{x_i}, D] = 0$. Then D is $\mathbb{R}[x_1, \dots, x_n]$ -linear. Since D vanishes on polynomials of degree k and is $\mathbb{R}[x_1, \dots, x_n]$ -linear, it vanishes on all polynomials.

Step 2: Let $f \in C^\infty \mathbb{R}^n$ and $x \in \mathbb{R}^n$ be a point. Using Hadamar's lemma, we obtain that $f = P + f_0$, where $P \in \mathbb{R}[x_1, \dots, x_n]$ is a polynomial of degree k and f_0 has zero of order $\geq k + 1$ in x . Then $D(f)(x) = D(P) + D(f_0) = 0$ (the first summand vanishes because D is $\mathbb{R}[x_1, \dots, x_n]$ -linear, and the second because D is of order k and f_0 has zero of order $\leq k + 1$ in x). ■

The ring of symbols

THEOREM: Consider the filtration $\text{Diff}^0(M) \subset \text{Diff}^1(M) \subset \text{Diff}^2(M) \subset \dots$ on the ring of differential operators. Then **its associated graded ring is isomorphic to the ring $\bigoplus_i \text{Sym}^i(TM)$.**

Proof. Step 1: Let f be a function with zero of order $\geq k$ in z , and \mathfrak{m} its maximal ideal. Then $\text{Diff}^{k-1}(f) = 0$. This gives a bilinear pairing

$$(\text{Diff}^k(M) / \text{Diff}^{k-1}(M)) \times (\mathfrak{m}^k) / (\mathfrak{m}^{k-1}) \longrightarrow \mathbb{R}$$

mapping $D \otimes f$ to $Df(0)$. Since $(\mathfrak{m}^k) / (\mathfrak{m}^{k-1}) = \text{Sym}^k(T_z M)^*$, this pairing, applied to all $z \in M$ gives a natural map $\text{Diff}^k(M) / \text{Diff}^{k-1}(M) \xrightarrow{\sigma} \text{Sym}^k(TM)$.

It remains to prove that σ is an isomorphism. Since this pairing is local, it suffices to prove that it is an isomorphism for $M = \mathbb{R}^n$.

Step 2: Let x_1, \dots, x_n be coordinates in $M = \mathbb{R}^n$, and $D \in \text{Diff}^k(M)$. Then $\text{Sym}^k(TM) = C^\infty M[\frac{d}{dx_1}, \frac{d}{dx_2}, \dots, \frac{d}{dx_n}]$. Consider a homogeneous differential

monomial $D = f \frac{\partial^k}{\partial x_{i_1} \dots \partial x_{i_k}}$. Then $\sigma(D) = f \frac{d}{dx_{i_1}} \frac{d}{dx_{i_2}} \dots \frac{d}{dx_{i_k}}$. **Therefore, σ is**

surjective.

Step 3: Let $D \in \text{Diff}^k(\mathbb{R}^n)$ be a differential operator and \underline{D} its class in $\text{Diff}^k(\mathbb{R}^n) / \text{Diff}^{k-1}(\mathbb{R}^n)$ such that $\sigma(\underline{D}) = 0$. Since σ is evaluation on polynomials, D vanishes on all homogeneous polynomials of degree k . By Lemma 1 above, $D = 0$. ■

Symbols

THEOREM: Consider the filtration $\text{Diff}^0(M) \subset \text{Diff}^1(M) \subset \text{Diff}^2(M) \subset \dots$. Then **its associated graded ring is isomorphic to $\bigoplus_i \text{Sym}^i(TM)$** , identified with the ring of fiberwise polynomial functions on T^*M .

COROLLARY: Let F, G be vector bundles, and $\text{Diff}^0(F, G) \subset \text{Diff}^1(F, G) \subset \text{Diff}^2(F, G)$ the corresponding spaces of differential operators. **Then**

$$\text{Diff}^i(F, G) / \text{Diff}^{i-1}(F, G) = \text{Sym}^i(TM) \otimes \text{Hom}(F, G),$$

where Sym^i denotes the symmetric power (symmetric part of the tensor power).

DEFINITION: Let F, G be vector bundles, and $D \in \text{Diff}^i(F, G)$ a differential operator. Consider its class in $\text{Diff}^i(F, G) / \text{Diff}^{i-1}(F, G)$ as a $\text{Hom}(F, G)$ -valued function on $T^*(M)$ (polynomial of order i on each cotangent space). This function is called **the symbol** of D .

EXERCISE: Let $D : B \rightarrow B \otimes \Lambda^1 M$ be a first order differential operator. **Prove that D is a connection if and only if its symbol is equal to the identity operator $\text{Id} \in \text{Hom}(\Lambda^1 M \otimes (\text{Hom}(B, B \otimes \Lambda^1 M)))$**

EXERCISE: Prove that the symbol of the Laplacian operator $\Delta : \Lambda^* M \rightarrow \Lambda^* M$ on a Riemannian manifold M at $\xi \in T^*M$ **is equal to $|\xi|^2 \text{Id}_{\Lambda^* M}$** .

Elliptic operators

DEFINITION: Let F, G be vector bundles of the same rank. A differential operator $D : F \rightarrow G$ is called **elliptic** if its symbol $\sigma(D) \in \text{Hom}(F, G) \otimes \text{Sym}^i(TM)$ is invertible at each non-zero $\xi \in T^*M$.

EXAMPLE: Consider an operator of second order on $C^\infty(\mathbb{R}^n)$,

$$D = f_0 + \sum_{i=1}^n f_i \frac{\partial}{\partial x_i} + \sum_{i,j=1}^n f_{ij} \frac{\partial^2}{\partial x_i \partial x_j}.$$

Then the symbol of D is $f_{ij} \frac{\partial^2}{\partial x_i \partial x_j}$; **it is elliptic if and only if the symmetric form $f_{ij} dx_i dx_j$ is positive or negative definite everywhere in \mathbb{R}^n .**

EXERCISE: Prove that symbol of D^* is a $\text{Hom}(F, G)$ -valued function on $T^*(M)$ which is Hermitian adjoint to $\text{symb}(D)$.

Elliptic operators: main properties

The rest of the slides today are introduction to the main results about elliptic operators; complete proofs for some of them will be given later.

THEOREM: (Elliptic regularity)

Let D be an elliptic operator with smooth coefficients on a manifold (not necessarily compact), and $f \in \ker D$. **Then D is smooth, and real analytic if coefficients of D were real analytic.**

THEOREM: Let D be an elliptic operator with smooth coefficients on a compact manifold. **Then its kernel is finite-dimensional.** If $D : F \rightarrow F$ is self-adjoint, then **D can be diagonalized in an appropriate orthonormal basis in the space of sections of F , and its eigenvalues are discrete.**

Elliptic operators of second order

Let $D = f_0 + \sum_{i=1}^n f_i \frac{\partial}{\partial x_i} + \sum_{i,j=1}^n f_{ij} \frac{\partial^2}{\partial x_i \partial x_j}$ be an elliptic operator of second order. For second order operators, **we always assume that the symbol $f_{ij} dx_i dx_j$ is positive definite.**

THEOREM: (E. Hopf's maximum principle)

Let D be a second order elliptic operator on a manifold (not necessarily compact) such that $D(const) = 0$, and f a solution of an equation $D(f) = 0$. Assume that f has a local maximum. **Then $f = const$.**

THEOREM: (Harnack inequality)

Let D be a second order elliptic operator on a manifold M (not necessarily compact), and $\Omega \subset M$ an open subset with compact closure. For any $f \in C^\infty(\Omega)$, denote by $\text{Var}_\Omega(f)$ the number $\sup_\Omega(f) - \inf_\Omega(f)$. **Then there exists a constant C , depending only on D , M and Ω , such that for any $f \in \ker D$, one has $\text{Var}_\Omega(f) < C$.**

COROLLARY: Any pointwise converging sequence of functions $f \in \ker D$ converges uniformly.

Proof: Indeed, by Harnack's inequality, the solutions of $D(f) = 0$ are uniformly continuous, hence the pointwise convergence implies uniform convergence. ■

Eberhard Hopf (1902-1983)



Fredholm operators

DEFINITION: Let F be a vector bundle on a compact manifold. The L_p^2 -**topology** on the space of sections of F is a topology defined by the quadratic form $|f|^2 = \sum_{i=0}^p \int_M |\nabla^i f|^2$, for some connection and scalar product on F and $\Lambda^1 M$.

EXERCISE: Prove that this topology is independent from the choice of a connection and a metric.

DEFINITION: A continuous operator $\psi : A \rightarrow B$ on topological vector spaces is called **Fredholm** if its kernel is finite-dimensional, and its image is closed, and has finite codimension.

THEOREM: Let $D : F \rightarrow G$ be an elliptic operator of order d . Clearly, D defines a continuous map $L_p^2(F) \rightarrow L_{p-d}^2(G)$. **Then this map is Fredholm.**

REMARK: This difficult theorem is a foundation of Hodge theory (and many other things).

Index of an elliptic operator

REMARK: Let $D : F \longrightarrow G$ be an elliptic operator of order d , and $L_p^2(F) \longrightarrow L_{p-d}^2(G)$ the corresponding maps on L_p^2 -spaces.

DEFINITION: **Index** of a Fredholm operator D is the number $\dim \ker D - \dim \operatorname{coker} D = \dim \ker D - \dim \ker D^*$

REMARK: Index of an elliptic operator $D : L_p^2(F) \longrightarrow L_{p-d}^2(G)$ a priori depends on p , however, by elliptic regularity, **all elements of $\ker D$ and $\ker D^*$ are smooth**, hence $\dim \ker D \Big|_{L_p^2(F)}$ is independent from p .

EXERCISE: Let F_t be a continuous family of Fredholm operators. **Prove that the index of F_t is constant in t .**

COROLLARY: Let D_t be a continuous family of elliptic operators of order k . **Then $\operatorname{ind} D_0 = \operatorname{ind} D_1$.**

COROLLARY: Index of an elliptic operator **is determined by its symbol.**

Proof: For any elliptic operators D_0, D_1 with the same symbol, the operator $D_t := tD_1 + (1-t)D_0$ has the same symbol, hence D_0 can be deformed to D_1 continuously and **gives a continuous family of Fredholm operators.** ■

Atiyah-Singer index theorem

REMARK: The index of an elliptic operator is clearly constant under continuous change of its symbol. Therefore, **index depends only on the homotopy class of its symbol**, which can be considered as a non-degenerate section of $\text{Hom}(F, G)$ over $T^*M \setminus 0$. Homotopy classes of such sections are described explicitly in terms of characteristic classes of F , G and the topological K-theory of M . **Atiyah-Singer index formula** expresses the index of an elliptic operator as a polynomial function of these topological invariants.

EXAMPLE: Let M be a compact manifold, and $D : C^\infty M \rightarrow C^\infty M$ elliptic operator of second order. **Then $\text{ind} D = 0$.**

Proof: Locally, we can write $D = f_0 + \sum_{i=1}^n f_i \frac{\partial}{\partial x_i} + \sum_{i,j=1}^n f_{ij} \frac{\partial^2}{\partial x_i \partial x_j}$. The Laplacian Δ associated with the metric tensor f_{ij} has the same symbol, hence it suffices to prove $\text{ind} \Delta = 0$. However, Δ is self-adjoint, hence $\dim \ker \Delta = \dim \text{coker} \Delta$. ■

COROLLARY: For any second order elliptic operator D on $C^\infty M$ with $D(\text{const}) = 0$, **one has $\dim \text{coker} D = 1$.**

Proof: By the strong maximum principle, all functions $\ker D$ are constant, hence $\dim \ker D = 1$. **Then $\dim \text{coker} D = 1$ by the index theorem.** ■