

# **Complex geometry**

## **lecture 7: Kähler metrics on homogeneous spaces**

Misha Verbitsky

**HSE, room 306, 16:20,**

**October 14, 2020**

## Homogeneous spaces

**DEFINITION:** A Lie group is a smooth manifold equipped with a group structure such that the group operations are smooth. Lie group  $G$  **acts on a manifold**  $M$  if the group action is given by the smooth map  $G \times M \longrightarrow M$ .

**DEFINITION:** Let  $G$  be a Lie group acting on a manifold  $M$  transitively. Then  $M$  is called **a homogeneous space**. For any  $x \in M$  the subgroup  $\text{St}_x(G) = \{g \in G \mid g(x) = x\}$  is called **stabilizer of a point**  $x$ , or **isotropy subgroup**.

**CLAIM:** For any homogeneous manifold  $M$  with transitive action of  $G$ , **one has**  $M = G/H$ , where  $H = \text{St}_x(G)$  is an isotropy subgroup.

**Proof:** The natural surjective map  $G \longrightarrow M$  putting  $g$  to  $g(x)$  identifies  $M$  with the space of conjugacy classes  $G/H$ . ■

**REMARK:** Let  $g(x) = y$ . Then  $\text{St}_x(G)^g = \text{St}_y(G)$ : **all the isotropy groups are conjugate**.

## Isotropy representation

**DEFINITION:** Let  $M = G/H$  be a homogeneous space,  $x \in M$  and  $\text{St}_x(G)$  the corresponding stabilizer group. The **isotropy representation** is the natural action of  $\text{St}_x(G)$  on  $T_xM$ .

**DEFINITION:** A bilinear symmetric form (or any tensor)  $\Phi$  on a homogeneous manifold  $M = G/H$  is called **invariant** if it is mapped to itself by all diffeomorphisms which come from  $g \in G$ .

**REMARK:** Let  $\Phi_x$  be an isotropy invariant tensor on  $T_xM$ , where  $M = G/H$  is a homogeneous space. For any  $y \in M$  obtained as  $y = g(x)$ , consider the form  $\Phi_y$  on  $T_yM$  obtained as  $\Phi_y := g^*(\Phi)$ . The choice of  $g$  is not unique, however, for another  $g' \in G$  which satisfies  $g'(x) = y$ , we have  $g = g'h$  where  $h \in \text{St}_x(G)$ . Since  $\Phi$  is  $h$ -invariant, **the tensor  $\Phi_y$  is independent from the choice of  $g$ .**

We proved

**THEOREM:** Let  $M = G/H$  be a homogeneous space and  $x \in M$  a point. Then  $G$ -invariant tensors on  $M = G/H$  **are in bijective correspondence with isotropy invariant tensors** on the vector space  $T_xM$ . ■

## Kähler manifolds

**DEFINITION:** A Riemannian metric  $g$  on an almost complex manifold  $M$  is called **Hermitian** if  $g(Ix, Iy) = g(x, y)$ . In this case,  $g(x, Iy) = g(Ix, I^2y) = -g(y, Ix)$ , hence  $\omega(x, y) := g(x, Iy)$  is skew-symmetric.

**REMARK:** Given any Riemannian metric  $g$  on an almost complex manifold, a **Hermitian metric  $h$  can be obtained as  $h = g + I(g)$ , where  $I(g)(x, y) = g(I(x), I(y))$ .**

**DEFINITION:** The differential form  $\omega \in \Lambda^{1,1}(M)$  is called **the Hermitian form** of  $(M, I, g)$ .

**REMARK:** It is  $U(1)$ -invariant, hence **of Hodge type (1,1)**.

**REMARK:** In the triple  $I, g, \omega$ , **each element can be recovered from the other two.**

**DEFINITION:** A complex Hermitian manifold  $(M, I, \omega)$  is called **Kähler** if  $d\omega = 0$ . The cohomology class  $[\omega] \in H^2(M)$  of a form  $\omega$  is called **the Kähler class** of  $M$ , and  $\omega$  **the Kähler form**.

## Erich Kähler



(Erich Kähler: 1990)

**16 January 1906 - 31 May 2000**

## Chez les Weil. André et Simone

André Weil: 6 May 1906 - 6 August 1998.



*"Simone et André à Penthievre, 1918-1919"*

## Representations acting transitively on a sphere

**THEOREM:** Let  $G$  be a group acting on a vector space  $V$ . Suppose that  $G$  acts transitively on a unit sphere  $\{x \in V \mid g(x) = 1\}$ . **Then a  $G$ -invariant bilinear symmetric form is unique up to a constant multiplier.**

**Proof. Step 1:** Since  $G$  preserves the sphere, which is a level set of the quadratic form  $g$ ,  $g$  is  $G$ -invariant.

**Step 2:** For any  $G$ -invariant quadratic form  $g'$ , the function  $x \rightarrow \frac{g'(x)}{g(x)}$  is constant on spheres and invariant under homothety, hence it is constant. ■

**EXERCISE:** Let  $V$  be a representation of  $G$ , and suppose  $G$  acts transitively on a sphere. **Prove that  $V$  is an irreducible representation.**

**EXERCISE:** Prove the **Schur lemma:** let  $V$  be an irreducible representation of  $G$  over  $\mathbb{R}$ , and  $g$  a  $G$ -invariant positive definite bilinear symmetric form. **Then any  $G$ -invariant bilinear symmetric form is proportional to  $g$ .**

## Fubini-Study form

**EXAMPLE:** Consider the natural action of the unitary group  $U(n+1)$  on  $\mathbb{C}P^n$ . The stabilizer of a point is  $U(n) \times U(1)$ .

**THEOREM:** There exists an  $U(n+1)$ -invariant Riemann form on  $\mathbb{C}P^n$ . Moreover, **such a form is unique up to a constant multiplier, and Kähler.**

**REMARK:** This Riemannian structure is called **the Fubini-Study metric**, and its Hermitian form **the Fubini-Study form**.

**Proof. Step 1:** To construct a  $U(n+1)$ -invariant Riemann form on  $\mathbb{C}P^n$ , we take a  $U(n)$ -invariant form on  $T_x\mathbb{C}P^n$  and apply Theorem 1. A  $U(n)$ -invariant form on  $T_x\mathbb{C}P^n$  exists, because it is a standard representation.

**Step 2:** Uniqueness follows because the isotropy group acts transitively on a sphere. ■

**CLAIM: The Fubini-Study form is closed,** and the corresponding metric is Kähler.

**Proof:** Let  $\omega$  be a Fubini-Study form. Then  $d\omega$  is an isotropy-invariant 3-form on  $T_x\mathbb{C}P^n$ . However, the isotropy group contains  $-\text{Id}$ , **hence all isotropy-invariant odd tensors vanish.** ■



## Projective manifolds

**DEFINITION:** Let  $M$  be a complex manifold, and  $X \subset M$  a smooth submanifold. It is called **a complex submanifold** if  $I(TX) \subset TX$ , and the map  $X \hookrightarrow M$  **a complex embedding**. A complex manifold which admits a complex embedding to  $\mathbb{C}P^n$  is called **a projective manifold**.

**REMARK:** **A complex submanifold of a Kähler manifold is Kähler.** Indeed, restriction of a Hermitian metric is Hermitian, and restriction of a closed form is closed. Therefore, **all projective manifolds are Kähler.**

**DEFINITION:** A subvariety of  $\mathbb{C}P^n$  is called **complex algebraic** if can be obtained as common zeroes of a system of homogeneous polynomial equations.

**THEOREM: (Chow theorem)** **All complex submanifolds in  $\mathbb{C}P^n$  are complex algebraic.**

## Kodaira embedding theorem

**DEFINITION: Kähler class** of a Kähler manifold is the cohomology class  $[\omega] \in H^2(M, \mathbb{R})$  of its Kähler form. We say that  $M$  **has integer Kähler class** if  $[\omega]$  belongs to the image of  $H^2(M, \mathbb{Z})$  in  $H^2(M, \mathbb{R})$

**REMARK:**  $H^2(\mathbb{C}P^n, \mathbb{R}) = \mathbb{R}$ . This implies that **the cohomology class of Fubini-Study form can be chosen integer**. In particular, **all projective manifolds admit Kähler structures with integer Kähler classes**.

**THEOREM: (Kodaira embedding theorem)** Let  $M$  be a compact Kähler manifold with an integer Kähler class. **Then it is projective**.

**This theorem will be proven later in these lectures.**

## Classes of almost complex manifolds

