

Brody lemma and Kobayashi hyperbolicity

Misha Verbitsky

November 7, Thursday, 2024,

seminar on geometric structures on manifolds, IMPA

Space forms

DEFINITION: **Simply connected space form** is a homogeneous manifold of one of the following types:

positive curvature: S^n (an n -dimensional sphere), equipped with an action of the group $SO(n+1)$ of rotations

zero curvature: \mathbb{R}^n (an n -dimensional Euclidean space), equipped with an action of isometries

negative curvature: $SO(1, n)/SO(n)$, equipped with the natural $SO(1, n)$ -action. This space is also called **hyperbolic space**, and in dimension 2 **hyperbolic plane** or **Poincaré plane** or **Bolyai-Lobachevsky plane**

LEMMA: Let $G = SO(n)$ act on \mathbb{R}^n in a natural way. **Then there exists a unique G -invariant symmetric 2-form:** the standard Euclidean metric.

COROLLARY: Let $M = G/H$ be a simply connected space form. **Then M admits a unique, up to a constant multiplier, G -invariant Riemannian form.**

Proof: The isotropy group is $SO(n-1)$ in all three cases, and the previous lemma can be applied. ■

REMARK: From now on, the space forms are assumed to be homogeneous Riemannian manifolds.

Schwartz lemma

CLAIM: (maximum principle) Let f be a holomorphic function defined on an open set U . **Then f cannot have strict maxima in U . If f has non-strict maxima, it is constant.**

EXERCISE: Prove the maximum principle.

LEMMA: (Schwartz lemma) Let $f : \Delta \rightarrow \Delta$ be a map from disk to itself fixing 0. **Then $|f'(0)| \leq 1$, and equality can be realized only if $f(z) = \alpha z$ for some $\alpha \in \mathbb{C}$, $|\alpha| = 1$.**

Proof: Consider the function $\varphi := \frac{f(z)}{z}$. Since $f(0) = 0$, it is holomorphic, and since $f(\Delta) \subset \Delta$, on the boundary $\partial\Delta$ we have $|\varphi|_{\partial\Delta} \leq 1$. Now, **the maximum principle implies that $|f'(0)| = |\varphi(0)| \leq 1$** , and equality is realized only if $\varphi = \text{const}$. ■

Conformal automorphisms of the disk

CLAIM: Let $\Delta \subset \mathbb{C}$ be the unit disk. **Then the group $\text{Aut}(\Delta)$ of its holomorphic automorphisms acts on Δ transitively.**

Proof: Let $V_a(z) = \frac{z-a}{1-\bar{a}z}$ for some $a \in \Delta$. Then $V_a(0) = -a$. To prove transitivity, it remains to show that $V_a(\Delta) = \Delta$, which is implied from

$$|V_a(z)| = |V_a(z)||z| = \left| \frac{z\bar{z} - a\bar{z}}{1 - \bar{a}z} \right| = \left| \frac{1 - a\bar{z}}{1 - \bar{a}z} \right| = 1.$$

■

REMARK: The group $PU(1, 1) \subset PGL(2, \mathbb{C})$ of unitary matrices preserving a pseudo-Hermitian form h of signature $(1, 1)$ acts on a disk $\{l \in \mathbb{C}P^1 \mid h(l, l) > 0\}$ by holomorphic automorphisms.

COROLLARY: Let $\Delta \subset \mathbb{C}$ be the unit disk, $\text{Aut}(\Delta)$ the group of its conformal automorphisms, and $\Psi : PU(1, 1) \longrightarrow \text{Aut}(\Delta)$ the map constructed above. **Then Ψ is an isomorphism.**

COROLLARY: Let h be a homogeneous metric on $\Delta = PU(1, 1)/S^1$. **Then (Δ, h) is conformally equivalent to $(\Delta, \text{flat metric})$.**

Upper half-plane

REMARK: The map $z \longrightarrow -\sqrt{-1}(z-1)^{-1}$ induces a diffeomorphism from the unit disc in \mathbb{C} to the upper half-plane \mathbb{H} .

PROPOSITION: The group $\text{Aut}(\Delta)$ acts on the upper half-plane \mathbb{H} as $z \xrightarrow{A} \frac{az+b}{cz+d}$, where $a, b, c, d \in \mathbb{R}$, and $\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} > 0$.

REMARK: The group of such A is naturally identified with $PSL(2, \mathbb{R}) \subset PSL(2, \mathbb{C})$. Since $PSL(2, \mathbb{R})$ acts on its Lie algebra preserving the Killing form, $PSL(2, \mathbb{R})$ embeds to $SO(1, 2)$. Both of these groups are 3-dimensional, since they are isomorphic.

REMARK: We have shown that $\mathbb{H} = SO(1, 2)/S^1$. This gives a **natural isomorphism of \mathbb{H} and the hyperbolic space**. Under this isomorphism, **holomorphic automorphisms correspond to isometries**.

Poincaré metric on disk

DEFINITION: Poincaré metric on a unit disk $\Delta \subset \mathbb{C}$ is an $\text{Aut}(\Delta)$ -invariant metric (it is unique up to a constant multiplier).

DEFINITION: Let $f : M \rightarrow M_1$ be a map of metric spaces. Then f is called **C -Lipschitz** if $d(x, y) \geq C d(f(x), f(y))$. A map is called **Lipschitz** if it is C -Lipschitz for some $C > 0$.

THEOREM: (Schwartz-Pick lemma)

Any holomorphic map $\varphi : \Delta \rightarrow \Delta$ from a unit disk to itself is 1-Lipschitz with respect to the Poincaré metric.

Proof. Step 1: We need to prove that for each $x \in \Delta$ the norm of the differential satisfies $|D\varphi_x| \leq 1$. Since the automorphism group acts on Δ transitively, **it suffices to prove that $|D\varphi_x| \leq 1$ when $x = 0$ and $\varphi(x) = 0$.**

Step 2: This is Schwartz lemma. ■

Kobayashi pseudometric

DEFINITION: Pseudometric on M is a function $d : M \times M \rightarrow \mathbb{R}^{\geq 0}$ which is symmetric: $d(x, y) = d(y, x)$ and satisfies the triangle inequality $d(x, y) + d(y, z) \geq d(x, z)$.

REMARK: Let \mathcal{D} be a set of pseudometrics. **Then** $d_{\max}(x, y) := \sup_{d \in \mathcal{D}} d(x, y)$ **is also a pseudometric.**

DEFINITION: The **Kobayashi pseudometric** on a complex manifold M is d_{\max} for the set \mathcal{D} of all pseudometrics such that any holomorphic map from the Poincaré disk to M is distance-decreasing.

EXERCISE: Prove that **the distance between points x, y in Kobayashi pseudometric is infimum of the Poincaré distance over all sets of Poincaré disks connecting x to y .**

EXAMPLE: The Kobayashi pseudometric on \mathbb{C} vanishes.

CLAIM: Any holomorphic map $X \xrightarrow{\varphi} Y$ is **1-Lipschitz with respect to the Kobayashi pseudometric.**

Proof: If $x \in X$ is connected to x' by a sequence of Poincaré disks $\Delta_1, \dots, \Delta_n$, then $\varphi(x)$ is connected to $\varphi(x')$ by $\varphi(\Delta_1), \dots, \varphi(\Delta_n)$. ■

Kobayashi hyperbolic manifolds

COROLLARY: Let $B \subset \mathbb{C}^n$ be a unit ball, and $x, y \in B$ points with coordinates $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n)$. Since x_i, y_i belongs to Δ , it makes sense to compute the Poincare distance $d_P(x_i, y_i)$. **Then $d_K(x, y) \geq \max_i d_P(x_i, y_i)$.**

Proof: Each of projection maps $\Pi_i : B \rightarrow \Delta$ is 1-Lipschitz. ■

DEFINITION: A variety is called **Kobayashi hyperbolic** if the Kobayashi pseudometric d_K is non-degenerate.

DEFINITION: A **domain** in \mathbb{C}^n is an open subset. A **bounded domain** is an open subset contained in a ball.

COROLLARY: **Any bounded domain Ω in \mathbb{C}^n is Kobayashi hyperbolic.**

Proof: Without restricting generality, we may assume that $\Omega \subset B$ where B is an open ball. Then the Kobayashi distance in Ω is \geq that in B . However, the Kobayashi distance in B is bounded by the metric $d(x, y) := \max_i d_P(x_i, y_i)$ as follows from above. ■

Caratheodory metric

DEFINITION: Let $x, y \in M$ be points on a complex manifold. Define **Caratheodory pseudometric** as $d_C(x, y) = \sup\{d_P(f(x), f(y))\}$, where the supremum is taken over all holomorphic map $f : M \rightarrow \Delta$, and d_P is Poincare metric on the disk Δ .

REMARK: Usually the term “Kobayashi/Caratheodory pseudometric” is abbreviated to “Kobayashi/Caratheodory metric”, **even when it is not a metric.**

REMARK: Caratheodory pseudometric **satisfies the triangle inequality** because a supremum of pseudometrics satisfies triangle inequality.

Exercise: Prove that **Caratheodory pseudometric is bounded by the Kobayashi pseudometric:** $d_K \geq d_C$.

REMARK: Clearly, $d_C \neq 0$ on any bounded domain.

Complex hyperbolic space

DEFINITION: Let $V = \mathbb{C}^{n+1}$ be a complex vector space equipped with a Hermitian metric h of signature $(1, n)$, and $\mathbb{H}_{\mathbb{C}}^n \subset \mathbb{P}V$ projectivization of the set of positive vectors $\{x \in V \mid h(x, \bar{x}) > 0\}$. Then $\mathbb{H}_{\mathbb{C}}^n$ is equipped with a homogeneous action of $U(1, n)$. The same argument as used for space forms implies that $\mathbb{H}_{\mathbb{C}}^n$ admits a $U(1, n)$ -invariant Hermitian metric, which is unique up to a constant multiplier. This Hermitian complex manifold is called **complex hyperbolic space**.

REMARK: For $n > 1$ it is not isometric, to the real hyperbolic spaces defined earlier.

REMARK: As a complex manifold $\mathbb{H}_{\mathbb{C}}^n$ is isomorphic to an open ball in \mathbb{C}^n .

REMARK: The Kobayashi metric and the Caratheodory metric on $\mathbb{H}_{\mathbb{C}}^n$ are $U(1, n)$ -invariant, because $U(1, n)$ acts holomorphically, hence proportional to the hyperbolic metric, which is also called **Bergman metric** on an open ball.

Exercise: Prove that **Kobayashi metric on a ball in \mathbb{C}^n is equal to the Caratheodory metric**.

Uniform convergence for Lipschitz maps

DEFINITION: A sequence of maps $f_i : M \rightarrow N$ between metric spaces **uniformly converges** (or **converges uniformly on compacts**) to $f : M \rightarrow N$ if for any compact $K \subset M$, we have $\lim_{i \rightarrow \infty} \sup_{x \in K} d(f_i(x), f(x)) = 0$.

Claim 1: Suppose that a sequence $f_i : M \rightarrow N$ of 1-Lipschitz maps converges to f pointwise in a countable dense subset $M' \subset M$. **Then f_i converges to f uniformly on compacts.**

Proof: Let $K \subset M$ be a compact set, and $N_\varepsilon \subset M'$ a finite subset such that K is a union of ε -balls centered in N_ε (such N_ε is called **an ε -net**). Then there exists N such that $\sup_{x \in N_\varepsilon} d(f_{N+i}(x), f(x)) < \varepsilon$ for all $i \geq 0$. Since f_i are 1-Lipschitz, this implies that

$$\begin{aligned} \sup_{y \in K} d(f_{N+i}(y), f(y)) &\leq \\ &\leq \sup_{x \in N_\varepsilon} d(f_{N+i}(x), f(x)) + \inf_{x \in N_\varepsilon} (d(f_{N+i}(x), y) + d(f(x), y)) \leq 3\varepsilon. \end{aligned}$$

■

Exercise: Prove that the limit f is also 1-Lipschitz.

REMARK: This proof works when M is a pseudo-metric space, as long as N is a metric space.

Arzelà-Ascoli theorem for Lipschitz maps

DEFINITION: Let M, N be metric spaces. A subset $B \subset M$ is **bounded** if it is contained in a ball. A family $\{f_\alpha\}$ of functions $f_\alpha : M \rightarrow N$ is called **uniformly bounded on compacts** if for any compact subset $K \subset M$, there is a bounded subset $C_K \subset N$ such that $f_\alpha(K) \subset C_K$ for any element f_α of the family.

THEOREM: (Arzelà-Ascoli for Lipschitz maps)

Let $\mathcal{F} := \{f_\alpha\}$ be an infinite uniformly bounded set of 1-Lipschitz maps $f_\alpha : M \rightarrow \mathbb{C}$, where M is a pseudo-metric space. Assume that M has countable base of open sets and can be obtained as a countable union of compact subsets. **Then there is a sequence $\{f_i\} \subset \mathcal{F}$ which converges to $f : M \rightarrow \mathbb{C}$ uniformly.**

REMARK: The limit f is clearly also 1-Lipschitz.

Proof. Step 1: Suppose we can prove Arzelà-Ascoli when M is compact. Then we can choose a sequence of compact subsets $K_i \subset M$, find subsequences in \mathcal{F} converging on each K_i , and use the diagonal method to find a subsequence converging on all K_i . Therefore, **we can assume that M is bounded, and all maps $f_\alpha : M \rightarrow \mathbb{C}$ map M into a compact subset $N \subset \mathbb{C}$.**

Arzelà-Ascoli theorem for Lipschitz maps (2)

THEOREM: (Arzelà-Ascoli for Lipschitz maps)

Let $\mathcal{F} := \{f_\alpha\}$ be an infinite uniformly bounded set of 1-Lipschitz maps $f_\alpha : M \rightarrow \mathbb{C}$, where M is a pseudo-metric space. Assume that M has countable base of open sets and can be obtained as a countable union of compact subsets. **Then there is a sequence $\{f_i\} \subset \mathcal{F}$ which converges to $f : M \rightarrow \mathbb{C}$ uniformly.**

REMARK: The limit f is clearly also 1-Lipschitz.

Proof. Step 1: We can assume that M is compact, and all maps $f_\alpha : M \rightarrow \mathbb{C}$ map M into a compact subset $N \subset \mathbb{C}$.

Step 2: Find a dense, countable subset $Z \subset M$. Using diagonal method, **find a sequence $\{f_i\} \subset \mathcal{F}$ converging pointwise to some f at all $z \in Z$.**

Step 3: Being a pointwise limit of Lipschitz functions, $f|_Z$ is also Lipschitz, and f_i converge to f uniformly on Z .

Step 4: Since a Lipschitz function maps Cauchy sequences to Cauchy sequences, it can be extended to a Lipschitz function on the completion \overline{M} .

■

Normal families of holomorphic functions

DEFINITION: Let M be a complex manifold. A family $\mathcal{F} := \{f_\alpha\}$ of holomorphic functions $f_\alpha : M \rightarrow \mathbb{C}$ is called **normal family** if \mathcal{F} is uniformly bounded on compact subsets.

THEOREM: (Montel's theorem)

Let M be a complex manifold with countable base, and \mathcal{F} a normal, infinite family of holomorphic functions. **Then there is a sequence $\{f_i\} \subset \mathcal{F}$ which converges to $f : M \rightarrow \mathbb{C}$ uniformly**, and f is holomorphic.

Proof. Step 1: As in the first step of Arzelà-Ascoli, it suffices to prove Montel's theorem on a subset of M where \mathcal{F} is bounded. Therefore, **we may assume that all f_α map M into a disk Δ .**

Step 2: All f_α are 1-Lipschitz with respect to Kobayashi metric. Therefore, **Arzelà-Ascoli theorem can be applied, giving a uniform limit $f = \lim f_i$.**

Step 3: A uniform limit of holomorphic functions is holomorphic by Cauchy formula. ■

REMARK: The sequence $f = \lim f_i$ **converges uniformly with all derivatives**, again by Cauchy formula.

Normal families in complete generality

DEFINITION: A set of holomorphic maps $f_\alpha : X \longrightarrow Y$ is called **a normal family** if any sequence $\{f_i\}$ in $\{f_\alpha\}$ has a subsequence converging uniformly on compacts.

THEOREM: Let $f_\alpha : X \longrightarrow Y$ be a family of holomorphic maps such that for any point $x \in X$ there exists its neighbourhood with compact closure $K \subset X$ and a Kobayashi hyperbolic open subset $V_K \subset Y$ such that all f_α map K to V_K . **Then f_α is a normal family.**

Proof: $f_\alpha|_K$ is Lipschitz with respect to the Kobayashi metric, and Arzelà-Ascoli theorem can be applied. ■

Brody curves and Brody maps

DEFINITION: Let M be a complex Hermitian manifold. **Brody curve** is a non-constant holomorphic map $f : \mathbb{C} \rightarrow M$ such that $|df| \leq C$ for some constant C . Here $|df|$ is understood as an operator norm of $df : T_z\mathbb{C} \rightarrow TM$, where \mathbb{C} is equipped with the standard Euclidean metric.

DEFINITION: Let (Δ_r, g_r) be a disk of radius r in \mathbb{C} with the Poincare metric g_r , rescaled in such a way that the unit tangent vector to 0 has length 1. **Brody map** to a Hermitian complex manifold is a map $f : \Delta_r \rightarrow M$ such that $|df| \leq 1$ (here the operator norm is taken with respect to the Poincare metric on Δ_r) and $|df|(z) = 1$ at $z = 0$.

Lemma 1: Let $f_r : \Delta_r \rightarrow M$ be a sequence of Brody maps with $r \rightarrow \infty$. **Then f_r converges uniformly to a Brody curve f satisfying $|df|(z) = 1$ at $z = 0$.**

Proof. Step 1: Let $r_1 < r_2$. **The identity map $\tau : (\Delta_{r_1}, g_{r_1}) \rightarrow (\Delta_{r_2}, g_{r_2})$ is 1-Lipschitz.** Indeed, it is 1-Lipschitz with respect to the usual Poincare metric: $\tau^*(r_2^{-2}g_{r_2}) \leq r_1^{-2}g_{r_1}$. Since $r_1 < r_2$, this gives $\tau^*g_{r_2} \leq g_{r_1}$.

Brody curves and Brody maps (2)

Lemma 1: Let $f_r : \Delta_r \rightarrow M$ be a sequence of Brody maps with $r \rightarrow \infty$. Then f_r converges uniformly to a Brody curve f satisfying $|df|(z) = 1$ at $z = 0$.

Proof. Step 1: Let $r_1 < r_2$. The identity map $\tau : (\Delta_{r_1}, g_{r_1}) \rightarrow (\Delta_{r_2}, g_{r_2})$ is 1-Lipschitz. Indeed, it is 1-Lipschitz with respect to the usual Poincare metric: $\tau^*(r_2^{-2}g_{r_2}) \leq r_1^{-2}g_{r_1}$. Since $r_1 < r_2$, this gives $\tau^*g_{r_2} \leq g_{r_1}$.

Step 2: Restricted on any disk Δ_R , the family $\{f_r, r > R\}$ is a normal family (it is Lipschitz), hence converges uniformly to a Lipschitz map. Since a uniform limit of holomorphic maps is holomorphic, the family $\{f_r|_{\Delta_R}, r > R\}$ converges to a holomorphic map on $\varphi_R : \Delta_R \rightarrow M$.

Step 3: The map φ_R is Lipschitz with respect to all metrics $g_r, r > R$. Since $\lim_r g_r$ is the standard Euclidean metric g_∞ , φ_R is Lipschitz with respect to g_∞ .

Step 4: $\lim_R \varphi_R$ converges to a holomorphic Lipschitz map $\mathbb{C} \rightarrow M$. Since all f_r and φ_R satisfy $|d\varphi_R|(z) = 1$ at $z = 0$, the same is true for the limit. ■

Brody curves and Brody maps

THEOREM: (Brody lemma)

Let M be a compact complex manifold which is not Kobayashi hyperbolic.

Then M contains a Brody curve.

Let us equip M with a Hermitian metric h . If $|df|(0) \leq C$ for any holomorphic map $(\Delta_1, g_1) \rightarrow M$, then the Kobayashi metric satisfies $d_K \geq C^{-1}h$, and M is Kobayashi hyperbolic. If it is non-bounded, we can always rescale the disc to obtain a map $f_r : (\Delta_r, g_r) \rightarrow M$ with $r = |df|(0)$, and then $|df_r|(0) = 1$.

Then Brody lemma follows from Lemma 1 and the following lemma.

LEMMA: Let M be a compact Hermitian manifold, and $\psi_r : (\Delta_r, g_r) \rightarrow M$ a sequence of holomorphic maps satisfying $|d\psi_r|(0) \geq 1$, $r \rightarrow \infty$. **Then there exists a sequence of Brody maps $f_s : (\Delta_s, g_s) \rightarrow M$, with $s \rightarrow \infty$.**

Brody curves and Brody maps (2)

LEMMA: Let M be a compact Hermitian manifold, and $\psi_r : (\Delta_r, g_r) \rightarrow M$ a sequence of holomorphic maps satisfying $|d\psi_r|(0) \geq 1$, $r \rightarrow \infty$. **Then there exists a sequence of Brody maps $f_s : (\Delta_s, g_s) \rightarrow M$, with $s \rightarrow \infty$.**

Proof. Step 1: We need to construct a sequence of holomorphic maps $f_s : (\Delta_s, g_s) \rightarrow M$, which are 1-Lipschitz and satisfy $|df_s|(0) = 1$. The identity map

$$\Psi_{r-\varepsilon, r} : (\Delta_{r-\varepsilon}, g_{r-\varepsilon}) \rightarrow (\Delta_r, g_r)$$

is 1-Lipschitz, and satisfies

$$\lim_{z \rightarrow \partial\Delta_{r-\varepsilon}} |d\Psi_{r-\varepsilon, r}|(z) = 0.$$

Let $u := r - \varepsilon$ and $\tilde{f}_u := \Psi_{r-\varepsilon, r} \circ \psi_r$ be a restriction of f_r to the disk $(\Delta_{r-\varepsilon}, g_{r-\varepsilon})$. Then \tilde{f}_u is also Lipschitz and $|d\tilde{f}_u|$ reaches maximum at a point z_u somewhere inside the disk Δ_u .

Step 2: Applying appropriate holomorphic isometry of Δ_u , we may assume that $|d\tilde{f}_u|(z)$ takes maximum $C_u \geq 1$ for $z = 0$. Rescaling \tilde{f}_u , and putting $s := C_u u$, we obtain a map $f_s : \Delta_s \rightarrow M$ which is 1-Lipschitz and satisfies $|df_s| \leq 1$, $|df_s|(0) = 1$. ■

Algebraically hyperbolic manifolds

DEFINITION: Let M be a projective manifold. We say that M is **algebraically hyperbolic** if there exists $A > 0$ such that for any curve $C \subset M$ of genus g one has $\deg C < A(g - 1)$.

REMARK: Algebraically hyperbolic manifolds **contain no elliptic nor rational curves.**

THEOREM: **Kobayashi hyperbolic implies algebraically hyperbolic.**

Converse implication (**“algebraically hyperbolic implies Kobayashi hyperbolic”**) was **conjectured by J.-P. Demailly** who introduced the notion of algebraic hyperbolicity.

Kobayashi hyperbolic implies algebraically hyperbolic

THEOREM: Kobayashi hyperbolic implies algebraically hyperbolic.

Proof. Step 1: Let $C \subset M$ be a curve in a Kobayashi hyperbolic manifold. Then its genus g is > 1 , and its universal covering is a disk. Denote by $\varphi_C : \Delta \rightarrow M$ the universal covering map. The volume of C with the Fubini-Study metric on M is $\deg C$, and its volume with the Poincare metric is $\int_C c_1(T^*C) = 2\pi(2g - 2)$. **Therefore, $\sup_{\Delta} |d\varphi_C| > \frac{\deg C}{2\pi(2g-2)}$.**

Step 2: Brody lemma implies that the Kobayashi metric d_K is bounded from below by εd_ω , where ω is a given Hermitian form on M , and d the corresponding distance function. Since any holomorphic map is Lipschitz with respect to the Kobayashi metric, this gives $\varepsilon d_\omega(\varphi_C(x), \varphi_C(y)) \leq d_K(\varphi_C(x), \varphi_C(y)) \leq d(x, y)$ for any $x, y \in \Delta$. **This gives $\varepsilon |d\varphi_C| \leq 1$.**

Step 3: Comparing Step 1 and Step 2, we obtain that $\varepsilon^{-1} \geq \sup_{\Delta} |d\varphi_C| > \frac{\deg C}{2\pi(2g-2)}$, hence $\frac{\deg C}{2\pi(2g-2)}$ is bounded from above. ■

Kobayashi hyperbolic implies algebraically hyperbolic: alternative proof

Another proof of the same statement

THEOREM: Kobayashi hyperbolic implies algebraically hyperbolic.

Proof. Step 1: Let $C \subset M$ be a curve in a Kobayashi hyperbolic manifold. Then its genus g is > 1 , and its universal cover is a disk. Denote by $\varphi_C : \Delta \rightarrow M$ the universal cover map. The volume of C with the Fubini-Study metric on M is $\deg C$, and its volume with Poincare metric is $\int_C c_1(T^*C) = 2\pi(2g - 2)$. **Therefore, $|\varphi_C| > \frac{\deg C}{2\pi(2g-2)}$ somewhere on Δ .**

Step 2: M is not algebraically hyperbolic \Leftrightarrow there is a sequence C_i of curves in M with $\lim_i \frac{\deg C_i}{2\pi(2g(C_i)-2)} \rightarrow \infty$. Suppose that M is not algebraically hyperbolic. Step 1 gives a sequence $\varphi_{C_i} : \Delta_1 \rightarrow M$ with $|\varphi_{C_i}| > \frac{\deg C_i}{2\pi(2g(C_i)-2)}$ somewhere on Δ . Replacing Δ_1 by $\Delta_{1-\varepsilon}$ as above, we may assume that $|\varphi_{C_i}|$ reaches maximum somewhere on Δ_1 . Applying isometry of Δ_1 , we may assume that $|\varphi_{C_i}|$ reaches maximum R_i in $0 \in \Delta_1$. Rescaling φ_{C_i} by R_i , we **obtain a sequence of disks $\tilde{\varphi}_{C_i}(z) = \varphi_{C_i}(z/R_i) : \Delta_{R_i} \rightarrow M$, giving a Brody curve (Lemma 1).** ■