

Ergodic complex structures and Kobayashi metric

Misha Verbitsky

Angers, Séminaire de géométrie algébrique

29/01/2014

Ergodic complex structures

DEFINITION: Let M be a smooth manifold. **A complex structure** on M is an endomorphism $I \in \text{End } TM$, $I^2 = -\text{Id}_{TM}$ such that the eigenspace bundles of I are **involutive**, that is, satisfy $[T^{1,0}M, T^{1,0}M] \subset T^{1,0}M$.

REMARK: Let Comp be the space of such tensors equipped with a topology of convergence of all derivatives. **It is a Fréchet manifold.**

DEFINITION: The diffeomorphism group Diff is a Fréchet Lie group acting on Comp in a natural way. A complex structure is called **ergodic** if its Diff -orbit is dense in Comp .

REMARK: The “moduli space” of complex structures (if it exists) is identified with $\text{Comp} / \text{Diff}$; **existence of ergodic complex structures guarantees that the moduli space does not exist.**

THEOREM: Let M be a compact torus, $\dim_{\mathbb{C}} M \geq 2$, or a simple hyperkähler manifold. **A complex structure on M is ergodic if and only if $\text{Pic}(M)$ is not of maximal rank.**

Teichmüller spaces

Definition: Let M be a compact complex manifold, and $\text{Diff}_0(M)$ a connected component of its diffeomorphism group (**the group of isotopies**). Denote by Comp the space of complex structures on M , and let $\text{Teich} := \text{Comp} / \text{Diff}_0(M)$. We call it **the Teichmüller space**.

Remark: Teich is **a finite-dimensional complex space** (Kodaira-Spencer-Kuranishi-Douady), but often **non-Hausdorff**.

Definition: Let $\text{Diff}_+(M)$ be the group of oriented diffeomorphisms of M . We call $\Gamma := \text{Diff}_+(M) / \text{Diff}_0(M)$ **the mapping class group**. The **coarse moduli space of complex structures on M** is a connected component of Teich / Γ .

REMARK: This terminology is **standard for curves**.

Moduli spaces

DEFINITION: The quotient $\text{Comp} / \text{Diff} = \text{Teich} / \Gamma$ is called **the moduli space** of complex structures. Typically, **it is very non-Hausdorff**. Comp corresponds bijectively to the set of isomorphism classes of complex structures.

REMARK: The moduli space exists, and is quasiprojective, for curves and manifolds with canonical polarization.

This talk is about an opposite situation, when Γ acts on Teich ergodically.

REMARK: Teich is a complex space for a wide class of manifolds, including all Calabi-Yau.

Holomorphically symplectic manifolds

DEFINITION: A **hyperkähler structure** on a manifold M is a Riemannian structure g and a triple of complex structures I, J, K , satisfying quaternionic relations $I \circ J = -J \circ I = K$, such that g is Kähler for I, J, K .

REMARK: A hyperkähler manifold is holomorphically symplectic: $\omega_J + \sqrt{-1} \omega_K$ is a holomorphic symplectic form on (M, I) .

THEOREM: (Calabi-Yau) A compact, Kähler, holomorphically symplectic manifold **admits a unique hyperkähler metric in any Kähler class.**

DEFINITION: For the rest of this talk, a hyperkähler manifold is a compact, Kähler, holomorphically symplectic manifold.

DEFINITION: A compact hyperkähler manifold M is called **simple** if $\pi_1(M) = 0$, $H^{2,0}(M) = \mathbb{C}$.

Bogomolov's decomposition: Any hyperkähler manifold admits a finite covering which is a product of a torus and several simple hyperkähler manifolds.

Further on, all hyperkähler manifolds are assumed to be simple.

Computation of the mapping class group

THEOREM: (Fujiki). Let $\eta \in H^2(M)$, and $\dim M = 2n$, where M is hyperkähler. **Then $\int_M \eta^{2n} = cq(\eta, \eta)^n$, for some primitive integer quadratic form q on $H^2(M, \mathbb{Z})$, and $c > 0$ a rational number.**

DEFINITION: The form q is called **Bogomolov-Beauville-Fujiki form**. It has signature $(3, b_2 - 3)$

THEOREM: (Sullivan) Let M be a compact, simply connected Kähler manifold, $\dim_{\mathbb{C}} M \geq 3$. Denote by Γ_0 the group of automorphisms of an algebra $H^*(M, \mathbb{Z})$ preserving the Pontryagin classes $p_i(M)$. Then **the natural map $\text{Diff}_+(M)/\text{Diff}_0 \rightarrow \Gamma_0$ has finite kernel, and its image has finite index in Γ_0 .**

THEOREM: Let M be a simple hyperkähler manifold, and Γ_0 as above. Then

- (i) $\Gamma_0|_{H^2(M, \mathbb{Z})}$ **is a finite index subgroup of $O(H^2(M, \mathbb{Z}), q)$.**
- (ii) The map $\Gamma_0 \rightarrow O(H^2(M, \mathbb{Z}), q)$ **has finite kernel.**

REMARK: Sullivan's theorem implies that the mapping class group for $\dim_{\mathbb{C}} M \geq 3$, $\pi_1(M) = 0$, **is an arithmetic lattice.**

The period map

Remark: For any $J \in \text{Teich}$, (M, J) is also a simple hyperkähler manifold, hence $H^{2,0}(M, J)$ is one-dimensional.

Definition: Let $P : \text{Teich} \rightarrow \mathbb{P}H^2(M, \mathbb{C})$ map J to a line $H^{2,0}(M, J) \in \mathbb{P}H^2(M, \mathbb{C})$. The map $P : \text{Teich} \rightarrow \mathbb{P}H^2(M, \mathbb{C})$ is called **the period map**.

REMARK: P maps Teich into an open subset of a quadric, defined by

$$\mathbb{P}er := \{l \in \mathbb{P}H^2(M, \mathbb{C}) \mid q(l, l) = 0, q(l, \bar{l}) > 0\}.$$

It is called **the period space** of M .

REMARK: $\mathbb{P}er = SO(b_2 - 3, 3)/SO(2) \times SO(b_2 - 3, 1)$. Indeed, the group $SO(H^2(M, \mathbb{R}), q) = SO(b_2 - 3, 3)$ acts transitively on $\mathbb{P}er$, and $SO(2) \times SO(b_2 - 3, 1)$ is a stabilizer of a point.

Birational Teichmüller moduli space

DEFINITION: Let M be a topological space. We say that $x, y \in M$ are **non-separable** (denoted by $x \sim y$) if for any open sets $V \ni x, U \ni y$, $U \cap V \neq \emptyset$.

THEOREM: (Huybrechts) Two points $I, I' \in \text{Teich}$ are **non-separable if and only if there exists a bimeromorphism $(M, I) \rightarrow (M, I')$ which is non-singular in codimension 2 and acts as identity on $H^2(M)$.**

REMARK: This is possible only if (M, I) and (M, I') contain a rational curve. **General hyperkähler manifold has no curves;** ones which have belong to a countable union of divisors in Teich .

DEFINITION: The space $\text{Teich}_b := \text{Teich} / \sim$ is called **the birational Teichmüller space** of M .

THEOREM: **The period map $\text{Teich}_b \xrightarrow{\text{Per}} \mathbb{P}\text{er}$ is an isomorphism,** for each connected component of Teich_b .

Ergodic complex structures

DEFINITION: Let (M, μ) be a space with measure, and G a group acting on M preserving measure. This action is **ergodic** if all G -invariant measurable subsets $M' \subset M$ satisfy $\mu(M') = 0$ or $\mu(M \setminus M') = 0$.

CLAIM: Let M be a manifold, μ a Lebesgue measure, and G a group acting on M ergodically. **Then the set of non-dense orbits has measure 0.**

Proof. Step 1: Consider a non-empty open subset $U \subset M$. Then $\mu(U) > 0$, hence $M' := G \cdot U$ satisfies $\mu(M \setminus M') = 0$. For any orbit $G \cdot x$ not intersecting U , $x \in M \setminus M'$. Therefore the set Z_U of such orbits has measure 0.

Proof. Step 2: Choose a countable base $\{U_i\}$ of topology on M . Then the set of points in dense orbits is $M \setminus \bigcup_i Z_{U_i}$. ■

DEFINITION: Let M be a complex manifold, Teich its Teichmüller space, and Γ the mapping group acting on Teich. **An ergodic complex structure** is a complex structure with dense Γ -orbit.

CLAIM: Let (M, I) be a manifold with ergodic complex structure, and I' another complex structure. **Then there exists a sequence of diffeomorphisms ν_i such that $\nu_i^*(I)$ converges to I' .**

Ergodicity of the monodromy group action

DEFINITION: A **lattice** in a Lie group is a discrete subgroup $\Gamma \subset G$ such that G/Γ has finite volume with respect to Haar measure.

THEOREM: (Calvin C. Moore, 1966) Let Γ be a lattice in a non-compact simple Lie group G with finite center, and $H \subset G$ a non-compact subgroup. **Then the left action of Γ on G/H is ergodic.**

THEOREM: Let $\mathbb{P}er$ be a component of a birational Teichmüller space, and Γ its monodromy group. Let $\mathbb{P}er_e$ be a set of all points $L \subset \mathbb{P}er$ such that the orbit $\Gamma \cdot L$ is dense (such points are called **ergodic**). **Then $Z := \mathbb{P}er \setminus \mathbb{P}er_e$ has measure 0.**

Proof. Step 1: Let $G = SO(b_2 - 3, 3)$, $H = SO(2) \times SO(b_2 - 3, 1)$. **Then Γ -action on G/H is ergodic,** by Moore's theorem.

Step 2: Ergodic orbits are dense, non-ergodic orbits have measure 0. ■

REMARK: Generic deformation of M has no rational curves, and no non-trivial birational models. Therefore, **outside of a measure zero subset,** $\text{Teich} = \text{Teich}_b$. This implies that **almost all complex structures on M are ergodic.**

Ratner's theorem

DEFINITION: Let G be a connected Lie group equipped with a Haar measure. **A lattice** $\Gamma \subset G$ is a discrete subgroup of finite covolume (that is, G/Γ has finite volume).

THEOREM: Let $H \subset G$ be a Lie subgroup generated by unipotents, and $\Gamma \subset G$ an arithmetic lattice. Then **a closure of any Γ -orbit in G/H is an orbit of a Lie subgroup $S \subset G$, such that $S \cap \Gamma \subset S$ is a lattice.**

EXAMPLE: Let V be a real vector space with a non-degenerate bilinear symmetric form of signature $(3, k)$, $k > 0$, $G := SO^+(V)$ a connected component of the isometry group, $H \subset G$ a subgroup fixing a given positive 2-dimensional plane, $H \cong SO^+(1, k) \times SO(2)$, and $\Gamma \subset G$ an arithmetic lattice. Consider the quotient $\mathbb{P}_{\text{er}} := G/H$. **Then a closure of $\Gamma \cdot J$ in G/H is an orbit of a closed connected Lie group $S \supset H$.**

Characterization of ergodic complex structures

CLAIM: Let $G = SO^+(3, k)$, and $H \cong SO^+(1, k) \times SO(2) \subset G$. Then **any closed connected Lie subgroup $S \subset G$ containing H coincides with G or with H .**

COROLLARY: Let $J \in \mathbb{P}er = G/H$. Then **either J is ergodic, or its Γ -orbit is closed in $\mathbb{P}er$.**

REMARK: By Ratner's theorem, in the latter case the H -orbit of J has finite volume in G/Γ . Therefore, **its intersection with Γ is a lattice in H .** This brings

COROLLARY: Let $J \in \mathbb{P}er$ be a point, such that its Γ -orbit is closed in $\mathbb{P}er$. Consider its stabilizer $St(J) \cong H \subset G$. **Then $St(J) \cap \Gamma$ is a lattice in $St(J)$.**

COROLLARY: Let J be a non-ergodic complex structure on a hyperkähler manifold, and $W \subset H^2(M, \mathbb{R})$ be a plane generated by $\operatorname{Re} \Omega, \operatorname{Im} \Omega$. **Then W is rational.**

REMARK: This can be used to show that **any hyperkähler manifold is Kobayashi non-hyperbolic.**

Kobayashi pseudometric

REMARK: The results further on are from a joint paper arXiv:1308.5667 by Ljudmila Kamenova, Steven Lu, Misha Verbitsky.

DEFINITION: Pseudometric on M is a function $d : M \times M \rightarrow \mathbb{R}^{\geq 0}$ which is symmetric: $d(x, y) = d(y, x)$ and satisfies the triangle inequality $d(x, y) + d(y, z) \geq d(x, z)$.

REMARK: Let \mathcal{D} be a set of pseudometrics. **Then** $d_{\max}(x, y) := \sup_{d \in \mathcal{D}} d(x, y)$ **is also a pseudometric.**

DEFINITION: The **Kobayashi pseudometric** on a complex manifold M is d_{\max} for the set \mathcal{D} of all pseudometrics such that any holomorphic map from the Poincaré disk to M is distance-decreasing.

THEOREM: Let $\pi : \mathcal{M} \rightarrow X$ be a smooth holomorphic family, which is trivialized as a smooth manifold: $\mathcal{M} = M \times X$, and d_x the Kobayashi metric on $\pi^{-1}(x)$. **Then** $d_x(m, m')$ **is upper continuous on x .** ■

COROLLARY: Denote the diameter of the Kobayashi pseudometric by $\text{diam}(d_x) := \sup_{m, m'} d_x(m, m')$. **Then** $\text{diam} : X \rightarrow \mathbb{R}^{\geq 0}$ **is upper continuous.**

Vanishing of Kobayashi pseudometric

THEOREM: Let (M, I) be a complex manifold with vanishing Kobayashi pseudometric. Then **the Kobayashi pseudometric vanishes for all ergodic complex structures in the same deformation class.**

Proof: Let $\text{diam} : \text{Comp} \rightarrow \mathbb{R}^{\geq 0}$ map a complex structure J to the diameter of the Kobayashi pseudometric on (M, J) . Let J be an ergodic complex structure. The set of points $J' = \nu(J) \in \text{Comp}$, $\nu \in \text{Diff}$, is dense, because J is ergodic. By upper semi-continuity, $0 = \text{diam}(I) \geq \inf_{J' = \nu(J)} \text{diam}(J') = \text{diam}(J)$.

■

EXAMPLE: Let M be a projective K3 surface. Then the Kobayashi metric on M vanishes. **Since all non-projective K3 are ergodic**, the Kobayashi metric vanishes on non-projective K3 surfaces as well.

THEOREM: Let M be a compact simple hyperkähler manifold. Assume that a deformation of M admits a holomorphic Lagrangian fibration and the Picard rank of M is not maximal. **Then the Kobayashi pseudometric on M vanishes.**

THEOREM: Let M be a Hilbert scheme of K3. **Then the Kobayashi pseudometric on M vanishes.**