

Global Torelli theorem for hyperkähler manifolds

Misha Verbitsky

Oberwolfach,

August 30, 2010.

Hyperkähler manifolds

DEFINITION: A **hyperkähler structure** on a manifold M is a Riemannian structure g and a triple of complex structures I, J, K , satisfying quaternionic relations $I \circ J = -J \circ I = K$, such that g is Kähler for I, J, K .

REMARK: A hyperkähler manifold **has three symplectic forms**
 $\omega_I := g(I\cdot, \cdot)$, $\omega_J := g(J\cdot, \cdot)$, $\omega_K := g(K\cdot, \cdot)$.

REMARK: This is equivalent to $\nabla I = \nabla J = \nabla K = 0$: the parallel translation along the connection preserves I, J, K .

DEFINITION: Let M be a Riemannian manifold, $x \in M$ a point. The subgroup of $GL(T_x M)$ generated by parallel translations (along all paths) is called **the holonomy group** of M .

REMARK: A hyperkähler manifold can be defined as a manifold which **has holonomy in $Sp(n)$** (the group of all endomorphisms preserving I, J, K).

Holomorphically symplectic manifolds

DEFINITION: A holomorphically symplectic manifold is a complex manifold equipped with non-degenerate, holomorphic $(2,0)$ -form.

REMARK: Hyperkähler manifolds are holomorphically symplectic. Indeed, $\Omega := \omega_J + \sqrt{-1} \omega_K$ is a holomorphic symplectic form on (M, I) .

THEOREM: (Calabi-Yau) A compact, Kähler, holomorphically symplectic manifold admits a unique hyperkähler metric in any Kähler class.

DEFINITION: For the rest of this talk, **a hyperkähler manifold is a compact, Kähler, holomorphically symplectic manifold.**

DEFINITION: A hyperkähler manifold M is called **simple** if $H^1(M) = 0$, $H^{2,0}(M) = \mathbb{C}$.

Bogomolov's decomposition: Any hyperkähler manifold admits a finite covering which is a product of a torus and several simple hyperkähler manifolds.

Remark: A simple hyperkähler manifold is always simply connected (Cheeger-Gromoll theorem).

Further on, all hyperkähler manifolds are assumed to be simple.

EXAMPLES.

EXAMPLE: An even-dimensional complex torus.

REMARK: Let M be a 2-dimensional complex manifold with holomorphic symplectic form outside of singularities, which are all of form $\mathbb{C}^2/\pm 1$. Then **its resolution is also holomorphically symplectic**. It is isomorphic to $T^*\mathbb{C}P^1$.

EXAMPLE: Take a 2-dimensional complex torus T , then all the singularities of $T/\pm 1$ are of this form. Its resolution $\widetilde{T/\pm 1}$ is called **a Kummer surface**. **It is holomorphically symplectic**.

REMARK: Take a symmetric square $\text{Sym}^2 T$, with a natural action of T , and let $T^{[2]}$ be a blow-up of a singular divisor. **Then $T^{[2]}$ is naturally isomorphic to the Kummer surface $\widetilde{T/\pm 1}$** .

DEFINITION: A **K3 surface** is a complex 2-manifold obtained as a deformation of a Kummer surface.

REMARK: **A K3 surface is always hyperkähler**. Any hyperkähler manifold of real dimension 4 is isomorphic to a torus or a K3 surface.

Hilbert schemes

DEFINITION: A **Hilbert scheme** $M^{[n]}$ of a complex surface M is a classifying space of all ideal sheaves $I \subset \mathcal{O}_M$ for which the quotient \mathcal{O}_M/I has dimension n over \mathbb{C} .

REMARK: A Hilbert scheme **is obtained as a resolution of singularities** of the symmetric power $\text{Sym}^n M$.

THEOREM: (Fujiki, Beauville) **A Hilbert scheme of a hyperkähler surface is hyperkähler.**

EXAMPLE: A Hilbert scheme of K3.

EXAMPLE: Let T is a torus. Then it acts on its Hilbert scheme freely and properly by translations. For $n = 2$, the quotient $T^{[n]}/T$ is a Kummer K3-surface. For $n > 2$, it is called **a generalized Kummer variety**.

REMARK: There are 2 more “sporadic” examples of compact hyperkähler manifolds, constructed by K. O’Grady. **All known simple compact hyperkähler manifolds are: Hilbert schemes of K3, generalized Kummer, and two O’Grady’s manifolds.**

Deformations of holomorphically symplectic manifolds.

REMARK: A Kaehler $2n$ -manifold (M, ω) with $c_1(M) = 0$ and a holomorphic 2-form Ω which satisfies $\int_M \Omega^n \wedge \bar{\Omega}^n \neq 0$ **is holomorphically symplectic.** Indeed, if Ω is degenerate at Z , Ω^n is a section of canonical class vanishing at $Z \subset M$. Then $c_1(M) = [Z]$, but $\langle [Z], \omega^{n-1} \rangle = \int_Z \omega^{n-1} > 0$.

COROLLARY: A small deformation of a holomorphically symplectic manifold **is again holomorphically symplectic.**

THEOREM: (Kodaira) **A small deformation of a compact Kähler manifold is again Kähler.**

COROLLARY: **Small deformations of hyperkähler manifolds are hyperkähler.**

REMARK: By **the moduli** of hyperkähler manifolds we shall understand the deformation space of complex manifolds admitting holomorphically symplectic and Kähler structure.

The Teichmüller space and the mapping class group

Definition: Let M be a compact complex manifold, and $\text{Diff}_0(M)$ a connected component of its diffeomorphism group (**the group of isotopies**). Denote by Teich the space of complex structures on M , and let $\text{Teich} := \text{Teich} / \text{Diff}_0(M)$. We call it **the Teichmüller space**.

Remark: Teich is a **finite-dimensional complex space** (Kodaira), but often **non-Hausdorff**.

Definition: Let $\text{Diff}_+(M)$ be the group of oriented diffeomorphisms of M . We call $\Gamma := \text{Diff}_+(M) / \text{Diff}_0(M)$ **the mapping class group**. The **coarse moduli space of complex structures on M** is a connected component of Teich / Γ .

Remark: This terminology is **standard for curves**.

REMARK: For hyperkähler manifolds, it is convenient to take for Teich **the space of all complex structures of hyperkähler type**, that is, **holomorphically symplectic and Kähler**. It is open in the usual Teichmüller space.

REMARK: To describe the moduli space, we shall compute Teich and Γ .

The Bogomolov-Beauville-Fujiki form

THEOREM: (Fujiki). Let $\eta \in H^2(M)$, and $\dim M = 2n$, where M is hyperkähler. Then $\int_M \eta^{2n} = cq(\eta, \eta)^n$, for some primitive integer quadratic form q on $H^2(M, \mathbb{Z})$, and $c > 0$ an integer number.

Definition: This form is called **Bogomolov-Beauville-Fujiki form**. It is defined by the Fujiki's relation uniquely, up to a sign. The sign is determined from the following formula (Bogomolov, Beauville)

$$\lambda q(\eta, \eta) = \int_X \eta \wedge \eta \wedge \Omega^{n-1} \wedge \overline{\Omega}^{n-1} - \frac{n-1}{n} \left(\int_X \eta \wedge \Omega^{n-1} \wedge \overline{\Omega}^n \right) \left(\int_X \eta \wedge \Omega^n \wedge \overline{\Omega}^{n-1} \right)$$

where Ω is the holomorphic symplectic form, and $\lambda > 0$.

Remark: q has signature $(b_2 - 3, 3)$. It is negative definite on primitive forms, and positive definite on $\langle \Omega, \overline{\Omega}, \omega \rangle$, where ω is a Kähler form.

Computation of the mapping class group

Theorem: (Sullivan) Let M be a compact simply connected Kähler manifold, $\dim_{\mathbb{C}} M \geq 3$. Denote by Γ_0 the group of automorphisms of an algebra $H^*(M, \mathbb{Z})$ preserving the Pontryagin classes $p_i(M)$. Then **the natural map $\text{Diff}_+(M)/\text{Diff}_0 \rightarrow \Gamma_0$ has finite kernel, and its image has finite index in Γ_0 .**

Theorem: Let M be a simple hyperkähler manifold, and Γ_0 as above. Then

- (i) $\Gamma_0|_{H^2(M, \mathbb{Z})}$ **is a finite index subgroup of $O(H^2(M, \mathbb{Z}), q)$.**
- (ii) The map $\Gamma_0 \rightarrow O(H^2(M, \mathbb{Z}), q)$ **has finite kernel.**

Proof. Step 1: Fujiki formula $v^{2n} = q(v, v)^n$ implies that Γ_0 **preserves the Bogomolov-Beauville-Fujiki up to a sign.** The sign is also fixed, if n is odd.

Step 2: For even n , the sign is also fixed. Indeed Γ_0 preserves $p_1(M)$, and (as Fujiki has shown) $v^{2n-2} \wedge p_1(M) = q(v, v)^{n-1}c$, for some $c \in \mathbb{R}$. The constant c is positive, **because the degree of $c_2(B)$ is positive** for any Yang-Mills bundle with $c_1(B) = 0$.

Computation of the mapping class group (cont.)

Step 3: $\mathfrak{o}(H^2(M, \mathbb{Q}), q)$ acts on $H^*(M, \mathbb{Q})$ by automorphisms preserving Pontryagin classes (V., 1995). Therefore $\Gamma_0|_{H^2(M, \mathbb{Q})}$ is an arithmetic subgroup of $O(H^2(M, \mathbb{R}), q)$.

Step 4: The kernel K of the map $\Gamma_0 \longrightarrow \Gamma_0|_{H^2(M, \mathbb{Q})}$ is finite, because it commutes with the Hodge decomposition and Lefschetz $\mathfrak{sl}(2)$ -action, hence preserves the Riemann-Hodge form, which is positive definite. ■

REMARK: The same argument as in Step 4 also proves that the group of automorphisms of $H^*(M, \mathbb{R})$ preserving p_1 is projected to $O(H^2(M, \mathbb{R}), q)$ with compact kernel.

REMARK: (Kollar-Matsusaka, Huybrechts) There are only finitely many connected components of Teich.

REMARK: The mapping class group acts on the set of connected components of Teich.

COROLLARY: Let Γ_I be the group of elements of mapping class group preserving a connected component of Teichmüller space containing $I \in \text{Teich}$. Then Γ_I is also arithmetic. Indeed, it has finite index in Γ .

The period map

Remark: For any $J \in \text{Teich}$, (M, J) is also a simple hyperkähler manifold, hence $H^{2,0}(M, J)$ is one-dimensional.

Definition: Let $P : \text{Teich} \rightarrow \mathbb{P}H^2(M, \mathbb{C})$ map J to a line $H^{2,0}(M, J) \in \mathbb{P}H^2(M, \mathbb{C})$. The map $P : \text{Teich} \rightarrow \mathbb{P}H^2(M, \mathbb{C})$ is called **the period map**.

REMARK: P maps Teich into an open subset of a quadric, defined by

$$\text{Per} := \{l \in \mathbb{P}H^2(M, \mathbb{C}) \mid q(l, l) = 0, q(l, \bar{l}) > 0\}.$$

It is called **the period space** of M .

REMARK: $\text{Per} = SO(b_2 - 3, 3)/SO(2) \times SO(b_2 - 3, 1)$

THEOREM: Let M be a simple hyperkähler manifold, and Teich its Teichmüller space. Then

- (i) (Bogomolov) **The period map $P : \text{Teich} \rightarrow \text{Per}$ is étale.**
- (ii) (Huybrechts) It is **surjective**.

REMARK: Bogomolov's theorem implies that Teich is smooth. It is **non-Hausdorff** even in the simplest examples.

Hausdorff reduction

REMARK: A non-Hausdorff manifold is a topological space locally diffeomorphic to \mathbb{R}^n .

DEFINITION: Let M be a topological space. We say that $x, y \in M$ are **non-separable** (denoted by $x \sim y$) if for any open sets $V \ni x, U \ni y, U \cap V \neq \emptyset$.

THEOREM: (D. Huybrechts) If $I_1, I_2 \in \text{Teich}$ are non-separable points, then $P(I_1) = P(I_2)$, and (M, I_1) **is birationally equivalent** to (M, I_2)

DEFINITION: Let M be a topological space for which M/\sim is Hausdorff. Then M/\sim is called **a Hausdorff reduction** of M .

Problems:

1. \sim **is not an equivalence relation.**
2. **Even if \sim is equivalence, the M/\sim is not always Hausdorff.**

REMARK: A quotient M/\sim is Hausdorff, if $M \rightarrow M/\sim$ is open, and the graph $\Gamma_{\sim} \in M \times M$ is closed.

Weakly Hausdorff manifolds

DEFINITION: A point $x \in X$ is called **Hausdorff** if $x \not\sim y$ for any $y \neq x$.

DEFINITION: Let M be an n -dimensional real analytic manifold, not necessarily Hausdorff. Suppose that **the set $Z \subset M$ of non-Hausdorff points is contained in a countable union of real analytic subvarieties of codim ≥ 2** . Suppose, moreover, that

(S) For every $x \in M$, there is a closed neighbourhood $B \subset M$ of x and a continuous surjective map $\Psi : B \rightarrow \mathbb{R}^n$ to a closed ball in \mathbb{R}^n , **inducing a homeomorphism** on an open neighbourhood of x .

Then M is called **a weakly Hausdorff manifold**.

REMARK: The period map satisfies (S). Also, the non-Hausdorff points of Teich **are contained in a countable union of divisors**.

THEOREM: A **weakly Hausdorff manifold X admits a Hausdorff reduction**. In other words, the quotient X/\sim is a Hausdorff. Moreover, $X \rightarrow X/\sim$ is locally a homeomorphism.

Birational Teichmüller moduli space

DEFINITION: The space $\text{Teich}_b := \text{Teich} / \sim$ is called **the birational Teichmüller space** of M .

THEOREM: **The period map** $\text{Teich}_b \xrightarrow{P} \mathbb{P}_{\text{er}}$ **is an isomorphism**, for each connected component of Teich_b .

Idea of a proof:

0. Since $\mathbb{P}_{\text{er}} = SO(b_2 - 3, 3) / SO(2) \times SO(b_2 - 3, 1)$ **is simply connected**, it will suffice to show it's a covering.
1. **It is etale** (Bogomolov).
2. For each hyperkähler structure (I, J, K) on M , **there is a whole S^2 of complex structures** $L = aI + bJ + cK$ on M , for $a^2 + b^2 + c^2 = 1$.
3. **On every such line, P is surjective.**

Birational Teichmüller moduli space (cont'd)

4. For any point $x \in \mathbb{P}er$, let $NS(x)$ be the corresponding lattice of integer $(1, 1)$ -classes in $H^2(M)$. Then **the set of Kaehler classes can be determined explicitly** when $\text{rk } NS(x) = 0$. Every Kaehler class gives a hyperkaehler structure, hence a line in Teich_b . Such a line is called **generic hyperkähler curve** (GHK curve).

5. **The whole Teichmüller space is covered by GHK curves.**

6. Surjectivity on GHK curves leads to the following condition.

(*) Let $\text{Teich}_b \xrightarrow{P} \mathbb{P}er$ be the period map. Then for each open subset $V \subset \mathbb{P}er$ with smooth boundary, and each connected component $W \subset P^{-1}(V)$, **the restriction $W \xrightarrow{P} V$ is surjective.**

6. The condition (*) **always implies that P is a covering.**

7. The period space is simply connected, hence **P is an isomorphism on each connected component.**

Global Torelli theorem

DEFINITION: Let M be a hyperkaehler manifold, Teich_b its birational Teichmüller space, and Γ the mapping class group. The quotient Teich_b/Γ is called **the birational moduli space** of M .

REMARK: The birational moduli space is obtained from the usual moduli space **by gluing some (but not all) non-separable points. It is still non-Hausdorff.**

THEOREM: Let (M, I) be a hyperkähler manifold, and W a connected component of its birational moduli space. **Then W is isomorphic to $\mathbb{P}er/\Gamma_I$, where $\mathbb{P}er = SO(b_2 - 3, 3)/SO(2) \times SO(b_2 - 3, 1)$ and Γ_I is an arithmetic group in $O(H^2(M, \mathbb{R}), q)$.**

A CAUTION: Usually “the global Torelli theorem” is understood as a theorem about Hodge structures. For K3 surfaces, **the Hodge structure on $H^2(M, \mathbb{Z})$ determines the complex structure.** For $\dim_{\mathbb{C}} M > 2$, **it is false.**

The Hodge-theoretic Torelli theorem

REMARK: The group $O(p, q)$ ($p, q > 0$) has **4 connected components**, corresponding to the orientations of positive p -dimensional and negative q -dimensional planes.

DEFINITION: Let M be a hyperkaehler manifold. One says that **the Hodge-theoretic Torelli theorem holds for M** if

$$\text{Teich} / \Gamma_I \longrightarrow \mathbb{P}er / O^+(H^2(M, \mathbb{Z}), q),$$

where $O^+(H^2(M, \mathbb{Z}), q)$ is a subgroup of $O(H^2(M, \mathbb{Z}), q)$ preserving orientation on positive 3-planes. Equivalently, **it is true if M is uniquely determined by its Hodge structure.**

REMARK: The Hodge-theoretic Torelli theorem **is true for K3 surfaces.** **It is false** for all other known examples of hyperkaehler manifolds.

Problems:

1. The moduli space Teich / Γ **is not Hausdorff** (Debarre, 1984). Indeed, bimeromorphically equivalent hyperkähler manifolds have isomorphic Hodge structures.
2. **The covering $\text{Teich}_b / \Gamma_I \longrightarrow \mathbb{P}er / O^+(H^2(M, \mathbb{Z}), q)$ is non-trivial**, because the map $\Gamma_I \longrightarrow O^+(H^2(M, \mathbb{Z}), q)$ is not surjective (Namikawa, 2002).

The birational Hodge-theoretic Torelli theorem

DEFINITION: The birational Hodge-theoretic Torelli theorem is true for M if Γ_I (the stabilizer of a Torelli component in the mapping class group) is isomorphic to $O^+(H^2(M, \mathbb{Z}), q)$.

REMARK: If a birational Hodge-theoretic Torelli theorem holds for M , then any deformation of M is up to a bimeromorphic equivalence **determined by the Hodge structure on $H^2(M)$** .

THEOREM: (Markman) The for $M = K3^{[n]}$, the group Γ_I **is a subgroup of $O^+(H^2(M, \mathbb{Z}), q)$ generated by oriented reflections**.

THEOREM: Let $M = K3^{[n+1]}$ with n a prime power. Then the (usual) global Torelli theorem holds birationally: **two deformations of a Hilbert scheme with isomorphic Hodge structures are bimeromorphic**. **For other n , it is false** (Markman).