

Holography principle and Moishezon twistor spaces

Misha Verbitsky

**Conference on The Geometry, Topology and Physics
of Moduli Spaces of Higgs Bundles**

National University of Singapore, 4-8 Aug 2014

Plan

1. Hyperkähler manifolds. Twistor spaces.
2. Holography principle
3. Moishezon twistor spaces. Applications of the holography principle to the local structure of twistor spaces.
4. Proof of holography principle.
- * 5. Hyperkähler reduction.

Questions

DEFINITION: A **hyperkähler structure** on a manifold M is a Riemannian structure g and a triple of complex structures I, J, K , satisfying quaternionic relations $I \circ J = -J \circ I = K$, such that g is Kähler for I, J, K .

REMARK: Any compact, Kähler, holomorphically symplectic manifold, such as a K3 surface, admits a hyperkähler metric, which is unique in its Kähler class.

QUESTION: Let (M, I, J, K) be a hyperkähler manifold. **Define when (M, I, J, K) is “algebraic”.**

REMARK: This question is motivated by [HKLR] definition of hyperkähler manifolds in terms of twistor spaces, which makes it possible to define a general “hyperkähler space” with singularities and nilpotents.

QUESTION: **Can an open ball in a K3 surface (or some other compact hyperkähler manifold) with its hyperkähler structure be equivalent to an open ball in a quiver space or a Hitchin moduli space with its hyperkähler structure?**

Twistor spaces

DEFINITION: Induced complex structures on a hyperkähler manifold are complex structures of form $L := aI + bJ + cK$. **It is a sphere:** $S^2 \cong \{L := aI + bJ + cK, \quad a^2 + b^2 + c^2 = 1.\}$

DEFINITION: A **twistor space** $\text{Tw}(M)$ of a hyperkähler manifold is a **complex manifold obtained by gluing these complex structures into a holomorphic family over $\mathbb{C}P^1$** . More formally:

Let $\text{Tw}(M) := M \times S^2$. Consider the complex structure $I_m : T_m M \rightarrow T_m M$ on M induced by $J \in S^2 \subset \mathbb{H}$. Let I_J denote the complex structure on $S^2 = \mathbb{C}P^1$.

The operator $I_{\text{Tw}} = I_m \oplus I_J : T_x \text{Tw}(M) \rightarrow T_x \text{Tw}(M)$ satisfies $I_{\text{Tw}}^2 = -\text{Id}$. **It defines an almost complex structure on $\text{Tw}(M)$** . This almost complex structure is known to be integrable (Obata)

Geometry of twistor spaces

EXAMPLE: If $M = \mathbb{H}^n$, $\text{Tw}(M) = \text{Tot}(\mathcal{O}(1)^{\oplus n}) \cong \mathbb{C}P^{2n+1} \setminus \mathbb{C}P^{2n-1}$ (total space of a vector bundle $(\mathcal{O}(1)^{\oplus n})$).

REMARK: (Deligne, Simpson)

The quaternionic structure on a hyperkähler manifold **can be reconstructed from the geometry of rational lines on its twistor space.**

REMARK: Twistor spaces can be defined for any conformally semiflat 4-dimensional Riemannian manifold M . These twistor space are often Moishezon (birational to projective). Results of today's talk ("holography principle") are true for such twistor spaces.

THEOREM: (Hitchin) For compact 4-dimensional M , its twistor space is Kähler only if $M = S^4$ or $M = \mathbb{C}P^2$.

Twistor spaces are non-Kähler

CLAIM: Suppose that (M, I, J, K) is a hyperkähler manifold such that (M, I) contains a compact, odd-dimensional complex subvariety Z . **Then $\text{Tw}(M)$ is non-Kähler.**

Proof: Consider the fundamental class $[Z_I] \in H_*(M)$ of Z in $(M, I) \subset \text{Tw}(M)$. Then $[Z_I] + [Z_{-I}] = 0$, giving $\int_{Z_I \cup Z_{-I}} \alpha = 0$ for each closed form $\alpha \in H^2(M)$. This is impossible if α is a Kähler form, because this integral is Riemannian volume of $Z_I \cup Z_{-I}$. ■

CLAIM: When M is compact and hyperkähler, $\text{Tw}(M)$ never admits a Kähler structure.

Proof: Let ω be the standard Hermitian form of $\text{Tw}(M)$. Then $dd^c\omega$ is a positive (2,2)-form (a calculation due to Kaledin-V.) For any Kähler form ω_0 , **this would imply**

$$\int_{\text{Tw}(M)} d\left(\omega_0^{\dim_{\mathbb{C}} M-1} \wedge d^c\omega\right) = \int_{\text{Tw}(M)} \omega_0^{\dim_{\mathbb{C}} M-1} \wedge dd^c\omega > 0,$$

which is impossible by Stokes' theorem. ■

Rational curves on $\text{Tw}(M)$.

DEFINITION: An ample rational curve on a complex manifold M is a smooth curve $S \cong \mathbb{C}P^1 \subset M$ such that $NS = \bigoplus_{k=1}^{n-1} \mathcal{O}(i_k)$, with $i_k > 0$. It is called a **quasiline** if all $i_k = 1$.

CLAIM: Let M be a compact complex manifold containing a an ample rational line. **Then any N points z_1, \dots, z_N can be connected by an ample rational curve.**

CLAIM: Let M be a hyperkähler manifold, $\text{Tw}(M) \xrightarrow{\sigma} M$ its twistor space, $m \in M$ a point, and $S_m = \mathbb{C}P^1 \times \{m\}$ the corresponding rational curve in $\text{Tw}(M)$. **Then S_m is a quasiline.**

Proof: Since the claim is essentially infinitesimal, it suffices to check it when M is flat. **Then $\text{Tw}(M) = \text{Tot}(\mathcal{O}(1)^{\oplus 2p}) \cong \mathbb{C}P^{2p+1} \setminus \mathbb{C}P^{2p-1}$, and S_m is a section of $\mathcal{O}(1)^{\oplus 2p}$. ■**

Holography principle

Theorem 1: (Holography principle for line bundles)

Let $S \subset M$ be an ample curve in a simply connected, connected complex manifold, which is covered by deformations of S . Consider a tubular neighbourhood $U \supset S$. Then, **for any holomorphic line bundle L on M , the space $H^0(U, L)$ is independent from the choice of U and S in its deformation class.**

REMARK: In these assumptions, $H^0(U, L)$ is always finite-dimensional (Hartshorne).

THEOREM: (Holography principle for meromorphic functions)

In assumptions of Theorem 1, **the space of meromorphic functions on U is equal to the space of meromorphic functions on M .**

DEFINITION: Given a complex manifold Z , denote by $\text{Mer}(Z)$ the field of global meromorphic functions on Z , and let **the algebraic dimension** $a(Z)$ be the transcendence degree of $\text{Mer}(Z)$.

REMARK: It could be infinite!

COROLLARY: $a(\text{Tw}(M)) = a(U)$ for any connected neighbourhood U of a quasiline in a connected, simply connected M .

Moishezon twistor spaces

DEFINITION: A compact complex variety Z is called **Moishezon** if the ring of meromorphic functions on Z has algebraic dimension $\dim Z$, or, equivalently, if Z is bimeromorphic to a projective manifold.

CLAIM: Let M be a simply connected hyperkaähler manifold, and $\text{Tw}(M)$ its twistor space. **Then** $a(\text{Tw}(M)) \leq \dim_{\mathbb{C}} \text{Tw}(M)$.

DEFINITION: A twistor space satisfying $a(\text{Tw}(M)) = \dim_{\mathbb{C}} \text{Tw}(M)$ is called **Moishezon**.

CLAIM: All Moishezon twistor spaces are bimeromorphic to open subsets of projective manifolds.

THEOREM: Let V be a quaternionic Hermitian vector space, and $G \subset \text{Sp}(V)$ a compact Lie group acting on V by quaternionic isometries. Denote by M the hyperkähler reduction of V . **Then $\text{Tw}(M)$ is Moishezon.**

Local structure of hyperkähler manifolds

THEOREM: (Fujiki, 1987) Let M be a compact hyperkähler manifold, and $L \in S^2 \subset \mathbb{H}$ a generic induced complex structure. **Then M contains no divisors.**

COROLLARY: Let M be a compact hyperkähler manifold, and $D \subset \text{Tw}(M)$ a divisor. **Then D is a union of several fibers of the projection $\pi : \text{Tw}(M) \rightarrow \mathbb{C}P^1$.**

Proof: Suppose that D contains a component which intersects generic fiber of π . By transversality, this intersection is a divisor, contradicting Fujiki's theorem. ■

COROLLARY: Let M be a compact, simply connected hyperkähler manifold, M' a hyperkähler manifold obtained by hyperkähler reduction (such as Nakajima quiver variety), and $U \subset M$, $U' \subset M'$ open subsets. **Then U is never equivalent to U' as a hyperkähler manifold.**

Proof: $a(\text{Tw}(U)) = a(\text{Tw}(M)) = 1$, and $a(\text{Tw}(U')) = a(\text{Tw}(M')) = \dim(\text{Tw}(M'))$. ■

Complex manifolds and quasilinear

REMARK: Let $S \subset M$ be a quasiline. Then, for an appropriate tubular neighbourhood $U \subset M$ of S , **“for every two points $x, y \in U$ close to S and far from each other, there is a unique deformation of S containing X and Y .”**

More precisely:

CLAIM: Let $S \subset M$ be a quasi-line. Then, for any sufficiently small tubular neighbourhood $U \subset M$ of S , there exists a smaller tubular neighbourhood $W \subset U$, satisfying the following condition. Let Δ_S be the image of the diagonal embedding $\Delta_S : S \rightarrow W \times W$. Then there exists an open neighbourhood V of Δ_S , properly contained in $W \times W$, such that **for any pair $(x, y) \in W \times W \setminus V$, there exists a unique deformation $S' \subset U$ of S containing x and y .**

COROLLARY: For any quasiline in M , its deformation space is $2(\dim M - 1)$ -dimensional.

Proof of holography principle

THEOREM: (Holography Principle)

Let $S \subset M$ be a quasiline in a simply connected complex manifold, which is covered by deformations of S . Assume that M is equipped with a projection $\pi : M \rightarrow S$ inducing identity on S . Consider a tubular neighbourhood $U \supset S$, and assume that $\pi : U \rightarrow S$ has connected fibers. Then, **for any holomorphic line bundle L on M , the space $H^0(U, L)$ is independent from the choice of U and S in its deformation class.**

A (slightly) weaker statement.

THEOREM 1: Let $S \subset M$ be a quasiline, and L a holomorphic bundle on M . Assume that M is equipped with a projection $\pi : M \rightarrow S$ inducing identity on S . Consider a sufficiently small tubular neighbourhood $U \supset S$, and a smaller tubular neighbourhood $V \subset U$. **Then the restriction map $H^0(U, L) \rightarrow H^0(V, L)$ is an isomorphism.**

We deduce the “Holography principle” from Theorem 1.

Deducing the holography principle from Theorem 1

Step 1: Choose a continuous, connected family S_b of quasilines parametrized by B such that $\cup_{b \in B} S_b = M$. Find a tubular neighbourhood U_b for each S_b in such a way that an intersection $U_b \cap U_{b'}$ for sufficiently close b, b' always contains S_b and $S_{b'}$. **By Theorem 1, $H^0(U_b \cap U_{b'}, L) = H^0(U_b, L) = H^0(U_{b'}, L)$.**

Step 2: Since B is connected, **all the spaces $H^0(U_b, L)$ are isomorphic**, and these isomorphisms are compatible with the restrictions to the intersections $U_b \cap U_{b'}$.

Step 3: Let now $f \in H^0(U_b, L)$, and let \tilde{M}_f be the **domain of holomorphy** for f , that is, a maximal domain (non-ramified over M) such that f admits a holomorphic extension to \tilde{M}_f . Since $\cup U_b = M$, and f can be holomorphically extended to any U_b , the domain \tilde{M}_f is a covering of M . **Now, “holography principle” follows, because M is simply connected. ■**

Proof of Theorem 1

THEOREM 1: Let $S \subset M$ be a quasiline, and L a holomorphic bundle on M . Assume that M is equipped with a projection $\pi : M \rightarrow S$ inducing identity on S . Consider a sufficiently small tubular neighbourhood $U \supset S$, and a smaller tubular neighbourhood $V \subset U$. **Then the restriction map $H^0(U, L) \rightarrow H^0(V, L)$ is an isomorphism.**

Proof. Step 1: Fix a point $x_0 = \infty$ in $\mathbb{C}P^1$. A section of $L|_{\mathbb{C}P^1} = \mathcal{O}(d)$ is the same as meromorphic function having a pole of degree $\leq d$ at $\pi^{-1}(\infty)$.

Step 2: For each section $S_1 : \mathbb{C}P^1 \rightarrow M$ of π , a degree d meromorphic function on S_1 is uniquely determined by its values at any $d + 1$ points of S_1 .

Step 3: Given a meromorphic function $f \in H^0(V, L)$ and a quasiline S_1 intersecting V in an open set, we can extend f to a meromorphic function \tilde{f} on S_1 by computing its values at $d + 1$ distinct points z_1, \dots, z_{d+1} of $S_1 \cap V$. Whenever S_1 is in V , this procedure gives $f|_{S_1}$. **By analytic continuation, the values of $\tilde{f}(z)$ at any $z \in S_1$ are independent from the choice of z_i and S_1 .**

Step 4: Therefore, \tilde{f} is a well-defined meromorphic function on the union V_1 of all quasilinear intersecting V . For U sufficiently small, V_1 contains U . Therefore, any $f \in H^0(V, L)$ can be extended to U . ■

Hamiltonians

Let's define the hyperkähler reduction.

We denote the Lie derivative along a vector field as $\text{Lie}_x : \Lambda^i M \longrightarrow \Lambda^i M$, and contraction with a vector field by $i_x : \Lambda^i M \longrightarrow \Lambda^{i-1} M$.

Cartan's formula: $d \circ i_x + i_x \circ d = \text{Lie}_x$.

REMARK: Let (M, ω) be a symplectic manifold, G a Lie group acting on M by symplectomorphisms, and \mathfrak{g} its Lie algebra. For any $g \in \mathfrak{g}$, denote by ρ_g the corresponding vector field. Then $\text{Lie}_{\rho_g} \omega = 0$, giving $d(i_{\rho_g}(\omega)) = 0$. **We obtain that $i_{\rho_g}(\omega)$ is closed, for any $g \in \mathfrak{g}$.**

DEFINITION: **A Hamiltonian** of $g \in \mathfrak{g}$ is a function h on M such that $dh = i_{\rho_g}(\omega)$.

Moment maps

DEFINITION: (M, ω) be a symplectic manifold, G a Lie group acting on M by symplectomorphisms. **A moment map** μ of this action is a linear map $\mathfrak{g} \rightarrow C^\infty M$ associating to each $g \in \mathfrak{g}$ its Hamiltonian.

REMARK: It is more convenient to consider μ as an element of $\mathfrak{g}^* \otimes_{\mathbb{R}} C^\infty M$, or (and this is most standard) **as a function with values in \mathfrak{g}^*** .

REMARK: Moment map **always exists** if M is simply connected.

DEFINITION: A moment map $M \rightarrow \mathfrak{g}^*$ is called **equivariant** if it is equivariant with respect to the coadjoint action of G on \mathfrak{g}^* .

REMARK: $M \xrightarrow{\mu} \mathfrak{g}^*$ is a moment map iff for all $g \in \mathfrak{g}$, $\langle d\mu, g \rangle = i_{\rho_g}(\omega)$. Therefore, **a moment map is defined up to a constant \mathfrak{g}^* -valued function**. An equivariant moment map is defined up to **a constant \mathfrak{g}^* -valued function which is G -invariant**.

DEFINITION: A G -invariant $c \in \mathfrak{g}^*$ is called **central**.

CLAIM: **An equivariant moment map exists whenever $H^1(G, \mathfrak{g}^*) = 0$** . In particular, when G is reductive and M is simply connected, an equivariant moment map exists.

Hyperkähler reduction

DEFINITION: Let G be a compact Lie group, ρ its action on a hyperkähler manifold M by hyperkähler isometries, and \mathfrak{g}^* a dual space to its Lie algebra. **A hyperkähler moment map** is a G -equivariant smooth map $\mu : M \rightarrow \mathfrak{g}^* \otimes \mathbb{R}^3$ such that $\langle \mu_i(v), g \rangle = \omega_i(v, d\rho(g))$, for every $v \in TM$, $g \in \mathfrak{g}$ and $i = 1, 2, 3$, where ω_i is one three Kähler forms associated with the hyperkähler structure.

DEFINITION: Let ξ_1, ξ_2, ξ_3 be three G -invariant vectors in \mathfrak{g}^* . The quotient manifold $M // G := \mu^{-1}(\xi_1, \xi_2, \xi_3) / G$ is called **the hyperkähler quotient** of M .

THEOREM: (Hitchin, Karlhede, Lindström, Roček)

The quotient $M // G$ is hyperkaehler.

Holomorphic moment map

Let $\Omega := \omega_J + \sqrt{-1}\omega_K$. This is a holomorphic symplectic (2,0)-form on (M, I) .

The proof of HKLR theorem. Step 1: Let μ_J, μ_K be the moment map associated with ω_J, ω_K , and $\mu_{\mathbb{C}} := \mu_J + \sqrt{-1}\mu_K$. Then $\langle d\mu_{\mathbb{C}}, g \rangle = i_{\rho g}(\Omega)$. Therefore, $d\mu_{\mathbb{C}} \in \Lambda^{1,0}(M, I) \otimes \mathfrak{g}^*$.

Step 2: This implies that the map $\mu_{\mathbb{C}}$ is holomorphic. It is called **the holomorphic moment map**.

Step 3: By definition, $M // G = \mu_{\mathbb{C}}^{-1}(c) // G$, where $c \in \mathfrak{g}^* \otimes_{\mathbb{R}} \mathbb{C}$ is a central element. **This is a Kähler manifold**, because it is a Kähler quotient of a Kähler manifold.

Step 4: We obtain 3 complex structures I, J, K on the hyperkähler quotient $M // G$. **They are compatible in the usual way** (an easy exercise). ■

Twistor spaces and hyperkähler reduction

THEOREM: Let V be a quaternionic Hermitian vector space, and $G \subset \mathrm{Sp}(V)$ a compact Lie group acting on V by quaternionic isometries. Denote by M the hyperkähler reduction of V . **Then $\mathrm{Tw}(M)$ is Moishezon.**

Proof: $\mathrm{Tw}(M)$ is obtained as the space of stable $G_{\mathbb{C}}$ -orbits in $\mu_{\mathbb{C}}^{-1}(0) \subset \mathrm{Tw}(V)$. The space $\mathrm{Tw}(V) = \mathbb{C}P^{2n+1} \setminus \mathbb{C}P^{2n-1}$ is algebraic. Averaging over G , we obtain that the field $G_{\mathbb{C}}$ -invariant rational functions on $\mathrm{Tw}(M)$ has dimension $\dim \mathrm{Tw}(M)$, and $\mathrm{Tw}(M)$ is Moishezon. ■

COROLLARY: Let U be an open subset of a compact, simply connected hyperkähler manifold, and U' an open subset of a hyperkähler manifold obtained as $V // G$, where V is flat and G reductive. **Then U is not isomorphic to U' as hyperkähler manifold.**

Proof: $\mathrm{Tw}(U')$ has many meromorphic functions, and $a(\mathrm{Tw}(U)) = 1$. ■